antibodies.com

## Datasheet for ABIN1694321 anti-C4BP antibody (AA 1-100) (AbBy Fluor® 350)



Overview

Quantity:	100 µL	
Target:	C4BP	
Binding Specificity:	AA 1-100	
Reactivity:	Human	
Host:	Rabbit	
Clonality:	Polyclonal	
Conjugate:	This C4BP antibody is conjugated to AbBy Fluor® 350	
Application:	Western Blotting (WB), Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence (Paraffin-embedded Sections) (IF (p))	

## Product Details

Immunogen:	KLH conjugated synthetic peptide derived from human C4 binding protein	
Isotype:	IgG	
Cross-Reactivity:	Human	
Predicted Reactivity:	Rat,Dog	
Purification:	Purified by Protein A.	
Target Details		
Target:	C4BP	
l arget:	СЧВР	

Alternative Name:	C4 Binding Protein (C4BP Products)	

Order at www.antibodies-online.com | www.antikoerper-online.de | www.anticorps-enligne.fr | www.antibodies-online.cn International: +49 (0)241 95 163 153 | USA & Canada: +1 877 302 8632 | support@antibodies-online.com Page 1/3 | Product datasheet for ABIN1694321 | 06/07/2024 | Copyright antibodies-online. All rights reserved.

Target Details	
Background:	Synonyms: C4b binding protein beta chain, C4b-binding protein beta chain, C4BP, C4BPB,
	C4BPB_HUMAN, Complement component 4 binding protein beta, Complement component 4
	binding protein beta chain, Complement component 4-binding protein beta,
	OTTHUMP00000034288, OTTHUMP00000034289, OTTHUMP00000034375,
	OTTHUMP0000003474, Proline rich protein, PRP.
	Background: C4BPB is a member of a superfamily of proteins composed predominantly of
	tandemly arrayed short consensus repeats of approximately 60 amino acids. A single, unique
	beta-chain encoded by this gene assembles with seven identical alpha-chains into the
	predominant isoform of C4b-binding protein, a multimeric protein that controls activation of the
	complement cascade through the classical pathway. C4b-binding protein has a regulatory role
	in the coagulation system also, mediated through the beta-chain binding of protein S, a vitamin
	K-dependent protein that serves as a cofactor of activated protein C. The genes encoding both
	alpha and beta chains are located adjacent to each other on human chromosome 1 in the
	regulator of complement activation gene cluster. Alternative splicing gives rise to multiple
	transcript variants

## Application Details

Application Notes:	IF(IHC-P) 1:50-200	
	IF(IHC-F) 1:50-200	
	IF(ICC) 1:50-200	
Restrictions:	For Research Use only	
Handling		
Format:	Liquid	
Concentration:	1 μg/μL	
Buffer:	Aqueous buffered solution containing 0.01M TBS ( pH 7.4) with 1 % BSA, 0.03 % Proclin300 and	
	50 % Glycerol.	
Preservative:	ProClin	
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be	
	handled by trained staff only.	
Storage:	-20 °C	
Storage Comment:	Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.	

Order at www.antibodies-online.com | www.antikoerper-online.de | www.anticorps-enligne.fr | www.antibodies-online.cn International: +49 (0)241 95 163 153 | USA & Canada: +1 877 302 8632 | support@antibodies-online.com Page 2/3 | Product datasheet for ABIN1694321 | 06/07/2024 | Copyright antibodies-online. All rights reserved.

1.1	(	1:
Н	land	ling
		3

Expiry Date:

12 months

Order at www.antibodies-online.com | www.antikoerper-online.de | www.anticorps-enligne.fr | www.antibodies-online.cn International: +49 (0)241 95 163 153 | USA & Canada: +1 877 302 8632 | support@antibodies-online.com Page 3/3 | Product datasheet for ABIN1694321 | 06/07/2024 | Copyright antibodies-online. All rights reserved.