antibodies .- online.com





anti-C1orf64 antibody (AA 81-169) (AbBy Fluor® 488)



Go to Product page

\sim			
	N/P	r\/I	i⊢₩

Quantity:	100 μL	
Target:	C1orf64 (C10RF64)	
Binding Specificity:	AA 81-169	
Reactivity:	Human	
Host:	Rabbit	
Clonality:	Polyclonal	
Conjugate:	This C1orf64 antibody is conjugated to AbBy Fluor® 488	
Application:	Western Blotting (WB), Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence (Paraffin-embedded Sections) (IF (p))	

Product Details

Immunogen:	KLH conjugated synthetic peptide derived from human C1orf64	
Isotype:	IgG	
Cross-Reactivity:	Human	
Purification:	Purified by Protein A.	

Target Details

Target:	C1orf64 (C1ORF64)
Alternative Name:	C1orf64 (C10RF64 Products)
Background: Synonyms: C1orf64, CA064_HUMAN, MGC24047, RP11-5P18.4, Uncharacterized protein	

C1orf64.

Background: Chromosome 1 is the largest human chromosome spanning about 260 million base pairs and making up 8 % of the human genome. There are about 3,000 genes on chromosome 1, and considering the great number of genes there are also a large number of diseases associated with chromosome 1. Notably, the rare aging disease Hutchinson-Gilford progeria is associated with the LMNA gene which encodes lamin A. When defective, the LMNA gene product can build up in the nucleus and cause characteristic nuclear blebs. The mechanism of rapidly enhanced aging is unclear and is a topic of continuing exploration. The MUTYH gene is located on chromosome 1 and is partially responsible for familial adenomatous polyposis. Stickler syndrome, Parkinsons, Gaucher disease and Usher syndrome are also associated with chromosome 1. A breakpoint has been identified in 1q which disrupts the DISC1 gene and is linked to schizophrenia. Aberrations in chromosome 1 are found in a variety of cancers including head and neck cancer, malignant melanoma and multiple myeloma. The C1orf64 gene product has been provisionally designated C1orf64 pending further characterization.

Gene ID:

149563

Application Details

Application	Notes:
, ipplication	1000.

IF(IHC-P) 1:50-200

IF(IHC-F) 1:50-200

IF(ICC) 1:50-200

Restrictions:

For Research Use only

Handling

Format:	Liquid
Concentration:	1 μg/μL
Buffer:	Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and 50 % Glycerol.
Preservative:	ProClin
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only.
Storage:	-20 °C

Handling

Storage Comment:	Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.
Expiry Date:	12 months