

[Go to Product page](#)

Datasheet for ABIN1694539

**anti-FUT6 antibody (AA 101-200) (Alexa Fluor 488)**

## Overview

Quantity:	100 µL
Target:	FUT6
Binding Specificity:	AA 101-200
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This FUT6 antibody is conjugated to Alexa Fluor 488
Application:	Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence (Paraffin-embedded Sections) (IF (p))

## Product Details

Immunogen:	KLH conjugated synthetic peptide derived from human FUT6/FucT-VI
Isotype:	IgG
Predicted Reactivity:	Human
Purification:	Purified by Protein A.

## Target Details

Target:	FUT6
Alternative Name:	FUT6 ( <a href="#">FUT6 Products</a> )
Background:	Synonyms: Alpha 1,3 fucosyltransferase, EC=2.4.1.65, FCT3A, Fuc TVI, Fucosyltransferase 6,

## Target Details

Fucosyltransferase VI, FucT VI, Galactoside 3 L fucosyltransferase, FUT6\_HUMAN.

Background: Glycosyltransferases that mediate the regio- and stereoselective transfer of sugars, such as the fucosyltransferases, determine cell surface-carbohydrate profiles, which is an essential interface for biological recognition processes. Fucosyltransferases catalyze the covalent association of fucose to different positional linkages in sugar acceptor molecules. The carbohydrate moieties generated and covalently attached to cell surfaces are necessary to ensure a surface contour that satisfies physiological roles, which are reliant on adhesion molecules such as Selectins (1-3). Hematopoietic lineages rely on Fucosyltransferases to confer a surface carbohydrate phenotype, which mediates proper cell adhesion molecule recruitment and cell trafficking (4-6).

Gene ID: 2528

## Application Details

Application Notes: IF(IHC-P) 1:50-200  
IF(IHC-F) 1:50-200  
IF(ICC) 1:50-200

Restrictions: For Research Use only

## Handling

Format: Liquid

Concentration: 1 µg/µL

Buffer: Aqueous buffered solution containing 0.01M TBS ( pH 7.4) with 1 % BSA, 0.03 % Proclin300 and 50 % Glycerol.

Preservative: ProClin

Precaution of Use: This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only.

Storage: -20 °C

Storage Comment: Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.

Expiry Date: 12 months