

Datasheet for ABIN1694577

anti-HGD antibody (AA 351-445) (AbBy Fluor® 488)[Go to Product page](#)

Overview

Quantity:	100 µL
Target:	HGD
Binding Specificity:	AA 351-445
Reactivity:	Mouse
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This HGD antibody is conjugated to AbBy Fluor® 488
Application:	Western Blotting (WB), Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence (Paraffin-embedded Sections) (IF (p))

Product Details

Immunogen:	KLH conjugated synthetic peptide derived from human HGD
Isotype:	IgG
Cross-Reactivity:	Mouse
Predicted Reactivity:	Human,Rat,Rabbit
Purification:	Purified by Protein A.

Target Details

Target:	HGD
Alternative Name:	Hgd (HGD Products)

Target Details

Background:	Synonyms: 2-dioxygenase, AKU, FLJ94126, hgd, HGD_HUMAN, HGO, Homogentisate 1 2 dioxygenase, Homogentisate 1, Homogentisate oxidase, Homogentisate oxygenase, Homogentisic acid oxidase, Homogentisicase. Background: HGD is a 445 amino acid protein that belongs to the homogentisate dioxygenase family and is involved in the pathway of amino acid degradation. Expressed at high levels in kidney, colon, liver, prostate and small intestine, HGD uses iron as a cofactor to catalyze the oxygen-dependent conversion of homogentisate to 4-maleylacetoacetate, a reaction that is the fourth step in the creation of L-phenylalanine from fumarate and acetoacetic acid. Defects in the gene encoding HGD are the cause of alkaptonuria (AKU), an autosomal recessive disorder that is characterized by urine that turns dark on standing and alkalinization, black ochronotic pigmentation of cartilage and collagenous tissues and spine arthritis.
-------------	--

Gene ID:	3081
----------	------

Application Details

Application Notes:	IF(IHC-P) 1:50-200 IF(IHC-F) 1:50-200 IF(ICC) 1:50-200
--------------------	--

Restrictions:	For Research Use only
---------------	-----------------------

Handling

Format:	Liquid
Concentration:	1 µg/µL
Buffer:	Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and 50 % Glycerol.
Preservative:	ProClin
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only.
Storage:	-20 °C
Storage Comment:	Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.
Expiry Date:	12 months