

# Datasheet for ABIN1694594

# anti-IgD antibody (AA 51-150) (AbBy Fluor® 488)



Overview	
Quantity:	100 μL
Target:	IgD
Binding Specificity:	AA 51-150
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This IgD antibody is conjugated to AbBy Fluor® 488
Application:	Western Blotting (WB), Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence (Paraffin-embedded Sections) (IF (p))
Product Details	
Immunogen:	KLH conjugated synthetic peptide derived from human IGHD
Isotype:	IgG
Predicted Reactivity:	Human
Purification:	Purified by Protein A.
Target Details	
Target:	IgD
Alternative Name:	Ighd (IgD Products)
Target Type:	Antibody

## **Target Details**

#### Background:

Synonyms: IGHD, Constant region of heavy chain of IgD, Ig delta chain C region, Ig delta chain C region secreted form, IgD, Igh-5, Immunoglobulin delta, Immunoglobulin heavy constant delta, IGHD\_HUMAN, Immunoglobulin heavy delta chain.

Background: Ig delta chain C region is an allelic product of the human IGHD gene. The two known IGHD alleles, IGHD\*01 and IGHD\*02, respectively produce isoforms 1, a secreted protein, and 2, a single-pass type I membrane protein. A member of the adaptive immune system, IgD are monomers expressed by activated B cells. Containing 3 Ig-like (immunoglobulin-like) domains, IgD chain C is located on chromosome 14 within the human heavy chain locus, lying on the 3? side of the IgM chain C region from the V-D-J cassette. Polyadenylation at certain sites along the heavy chain locus likely effects the mechanism that determines the alternative splicing event which results in the expression of either IgD chain C or IgM chain C. Some studies have suggested that antigenic coactivation of IgD+ B cells can have a negative influence on bone resorption during infectious events.

# **Application Details**

Application Notes:	IF(IHC-P) 1:50-200
	IF(IHC-F) 1:50-200
	IF(ICC) 1:50-200
Restrictions:	For Research Use only

## Handling

Format:	Liquid
Concentration:	1 μg/μL
Buffer:	Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and 50 % Glycerol.
Preservative:	ProClin
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only.
Storage:	-20 °C
Storage Comment:	Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.
Expiry Date:	12 months