antibodies .- online.com





anti-Acetyl-CoA Carboxylase beta antibody (pSer220) (AbBy Fluor® 488)



Go to Product page

()	11/	IN	/ie	A .
	/ // 	۱ ات	/ (−	' \/\/

Quantity:	100 μL	
Target:	Acetyl-CoA Carboxylase beta (ACACB)	
Binding Specificity:	pSer220	
Reactivity:	Human	
Host:	Rabbit	
Clonality:	Polyclonal	
Conjugate:	This Acetyl-CoA Carboxylase beta antibody is conjugated to AbBy Fluor® 488	
Application:	Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence (Paraffin-embedded Sections) (IF (p))	
Product Details		
Immunogen:	KLH conjugated synthetic phosphopeptide derived from human Acetyl Coenzyme A	
	Carboxylase beta around the phosphorylation site of Ser220	
Isotype:	IgG	
Predicted Reactivity:	Human	
Purification:	Purified by Protein A.	
Target Details		
Target:	Acetyl-CoA Carboxylase beta (ACACB)	
	Acetyl Coenzyme A Carboxylase beta (ACACB Products)	

Target Details

Storage Comment:

rarget Details		
Background:	Synonyms: Acetyl Coenzyme A Carboxylase beta phospho S220, Acetyl Coenzyme A Carboxylase beta phospho Ser2210 p-Acetyl Coenzyme A Carboxylase beta phospho S220, ACACB_HUMAN Background: Acetyl-CoA carboxylase (ACC) is a complex multifunctional enzyme system. ACC is a biotin-containing enzyme which catalyzes the carboxylation of acetyl-CoA to malonyl-CoA, the rate-limiting step in fatty acid synthesis. There are two ACC forms, alpha and beta, encoded by two different genes. ACC-alpha is highly enriched in lipogenic tissues. The enzyme is under long term control at the transcriptional and translational levels and under short term regulation by the phosphorylation/dephosphorylation of targeted serine residues and by allosteric transformation by citrate or palmitoyl-CoA. Multiple alternatively spliced transcript variants divergent in the 5' sequence and encoding distinct isoforms have been found for this gene.	
	[provided by RefSeq, Jul 2008].	
UniProt:	000763	
Pathways:	AMPK Signaling, Ribonucleoside Biosynthetic Process	
Application Details		
Application Notes:	IF(IHC-P) 1:50-200	
	IF(IHC-F) 1:50-200	
	IF(ICC) 1:50-200	
Restrictions:	For Research Use only	
Handling		
Format:	Liquid	
Concentration:	1 μg/μL	
Buffer:	Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and 50 % Glycerol.	
Preservative:	ProClin	
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only.	
Storage:	-20 °C	

Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.

\vdash	land	lına
	iaria	шц

Expiry Date:

12 months