# antibodies .- online.com





# anti-Allergin-1 antibody (AA 61-160) (Alexa Fluor 488)



Go to Product page

( )	1/0	r\ /1	014	
( )	ve	I V I	-v	V

Quantity:	100 μL	
Target:	Allergin-1 (MILR1)	
Binding Specificity:	AA 61-160	
Reactivity:	Human	
Host:	Rabbit	
Clonality:	Polyclonal	
Conjugate:	This Allergin-1 antibody is conjugated to Alexa Fluor 488	
Application:	Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence (Paraffin-embedded Sections) (IF (p))	

#### **Product Details**

Immunogen:	KLH conjugated synthetic peptide derived from human Allergin-1	
Isotype:	IgG	
Predicted Reactivity:	Human	
Purification:	Purified by Protein A.	

## **Target Details**

Target:	Allergin-1 (MILR1)	
Alternative Name:	Allergin-1 (MILR1 Products)	
Background:	Synonyms: C17orf60,Chromosome 17 open reading frame 60.	

Background: As a major event in type allergic reaction, degranulation of mast cells is triggered by the interaction between specific IgEs and high affinity IgE receptors (FcRIs) on mast cell membrane, and is followed by the release of a battery of inflammatory mediators, such as histamine, leukotrienes and arachidonic acid, and these inflammatory mediators in turn induce a series of allergic symptoms. Recently, Allergin-1 has been found to be an immunoglobulin-like receptor preferentially expressed on mast cells. The crosslinking of Allergin-1 to FcRIs on mast cells may inhibit the signal transduction from FcRIs to intracellular immunoreceptor tyrosine-based inhibitory motif (ITIM), generate ITIM inhibitory signals, which then participate in the regulation of mast cell degranulation, and finally block the initiation of allergic inflammation.

### **Application Details**

Application Notes:	IF(IHC-P) 1:50-200
	IF(IHC-F) 1:50-200
	IF(ICC) 1:50-200

Restrictions: For Research Use only

#### Handling

Format:	Liquid
Concentration:	1 μg/μL
Buffer:	Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and 50 % Glycerol.
Preservative:	ProClin
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only.
Storage:	-20 °C
Storage Comment:	Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.
Expiry Date:	12 months