# antibodies -online.com





## anti-TDRP/C8orf42 antibody (AA 51-150) (Alexa Fluor 488)



Go to Product page

$\sim$					
	1/6	⊃r	1//	$\triangle$	٨/

Quantity:	100 μL
Target:	TDRP/C8orf42 (TDRP)
Binding Specificity:	AA 51-150
Reactivity:	Rat
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This TDRP/C8orf42 antibody is conjugated to Alexa Fluor 488
Application:	Western Blotting (WB), Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence (Paraffin-embedded Sections) (IF (p))

#### **Product Details**

Immunogen:	KLH conjugated synthetic peptide derived from human C8orf42	
Isotype:	IgG	
Cross-Reactivity:	Rat	
Predicted Reactivity:	Human, Mouse	
Purification:	Purified by Protein A.	

#### **Target Details**

Target:	TDRP/C8orf42 (TDRP)
Alternative Name:	C8orf42 (TDRP Products)

### **Target Details**

Expiry Date:

Background:	Synonyms: C8orf42, TDRP_HUMAN, Protein INM01, Protein TDRP, Uncharacterized protein			
	C8orf42.			
	Background: Made up of nearly 146 million bases, chromosome 8 encodes about 800 genes.			
	Translocation of portions of chromosome 8 with amplifications of the c-Myc gene are found in			
	some leukemias and lymphomas, and typically associated with a poor prognosis. Portions of			
	chromosome 8 have been linked to schizophrenia and bipolar disorder. Trisomy 8, also known			
	as Warkany syndrome 2, most often results in early miscarriage but is occasionally seen in a			
	mosaic form in surviving patients who suffer to a varying degree from a number of symptoms			
	including retarded mental and motor development, and certain facial and developmental			
	defects. WRN is a DNA helicase encoded by chromosome 8 and shown defective in those with			
	the early aging disorder Werner syndrome. Chromosome 8 is also associated with Pfeiffer			
	syndrome, congenital hypothyroidism and Waardenburg syndrome. The C8orf42 gene product			
	has been provisionally designated C8orf42 pending further characterization.			
Gene ID:	157695			
Application Details				
Application Notes:	IF(IHC-P) 1:50-200			
	IF(IHC-F) 1:50-200			
	IF(ICC) 1:50-200			
Restrictions:	For Research Use only			
Handling				
Format:	Liquid			
Concentration:	1 μg/μL			
Buffer:	Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and			
	50 % Glycerol.			
Preservative:	ProClin			
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be			
	handled by trained staff only.			
Storage:	-20 °C			
Storage Comment:	Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.			
Evniny Data:	12 months			

12 months