# antibodies -online.com





## anti-Cadherin 12 antibody (AA 291-390) (AbBy Fluor® 488)



Go to Product page

$\sim$					
	1/6	⊃r	\/I	$\triangle$	٨/

Quantity:	100 μL	
Target:	Cadherin 12	
Binding Specificity:	AA 291-390	
Reactivity:	Rat	
Host:	Rabbit	
Clonality:	Polyclonal	
Conjugate:	This Cadherin 12 antibody is conjugated to AbBy Fluor® 488	
Application:	Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence (Paraffin-embedded Sections) (IF (p))	

#### **Product Details**

Immunogen:	KLH conjugated synthetic peptide derived from human CDH12	
Isotype:	IgG	
Cross-Reactivity:	Rat	
Predicted Reactivity:	Human, Mouse, Dog, Cow, Sheep, Pig, Horse, Chicken, Rabbit	
Purification:	Purified by Protein A.	

### **Target Details**

Target:	Cadherin 12	
Alternative Name:	Cdh12 (Cadherin 12 Products)	

#### **Target Details**

#### Background:

Synonyms: Br cadherin, BR-cadherin, Brain cadherin, CAD12\_HUMAN, cadherin 12 N cadherin 2, Cadherin 12, Cadherin 12 precursor, Cadherin 12 type 2 N cadherin 2, Cadherin 12 type 2, Cadherin 12, CDH12, CDHB, FLJ34857, N cadherin 2, N-cadherin 2, Neural type cadherin 2, Neuronal cadherin 2.

Background: The cadherins are a family of Ca++-dependent adhesion molecules that function to mediate cell-cell binding critical to the maintenance of tissue structure and morpho-genesis (1-4). Cadherins each contain a large extracellular domain at the amino terminus, which is characterized by a series of five homologous repeats, the most distal of which is thought to be responsible for binding specificity (5,6). The relatively short carboxy terminal, intracellular domain interacts with a variety of cytoplasmic proteins, including catenin b, to regulate cadherin function (7). BR-cadherin (for brain-cadherin, also designated cadherin-12 or N-cadherin 2) is expressed specifically in neurons in the central nervous system and is thought to be involved in neuronal development (8).

#### **Application Details**

Application Notes:	IF(IHC-P) 1:50-200
	IF(IHC-F) 1:50-200
	IF(ICC) 1:50-200
Restrictions:	For Research Use only

#### Handling

Format:	Liquid	
Concentration:	1 μg/μL	
Buffer:	Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and 50 % Glycerol.	
Preservative:	ProClin	
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only.	
Storage:	-20 °C	
Storage Comment:	Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.	
Expiry Date:	12 months	