antibodies .- online.com







anti-RHOG antibody (AA 61-150) (Alexa Fluor 488)



| () | 11/0 | r\ /1 | $\triangle 1 $ |
|-----|--------------------|-------|----------------|
| | $\lor \lor \vdash$ | 1 V I | ew |
| | | | |

| Quantity: | 100 μL |
|----------------------|--|
| Target: | RHOG |
| Binding Specificity: | AA 61-150 |
| Reactivity: | Human |
| Host: | Rabbit |
| Clonality: | Polyclonal |
| Conjugate: | This RHOG antibody is conjugated to Alexa Fluor 488 |
| Application: | Flow Cytometry (FACS), Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence (Paraffin-embedded Sections) (IF (p)) |

Product Details

| Immunogen: | KLH conjugated synthetic peptide derived from human RHOG |
|-----------------------|--|
| Isotype: | IgG |
| Cross-Reactivity: | Human |
| Predicted Reactivity: | Mouse,Rat |
| Purification: | Purified by Protein A. |

Target Details

| Target: | RHOG |
|-------------------|----------------------|
| Alternative Name: | RHOG (RHOG Products) |

Target Details

| Background: | Synonyms: ARHG, Rho-related GTP-binding protein RhoG, RHOG | |
|---------------------|--|--|
| | Background: Required for the formation of membrane ruffles during macropinocytosis. Plays a | |
| | role in cell migration and is required for the formation of cup-like structures during trans- | |
| | endothelial migration of leukocytes. In case of Salmonella enterica infection, activated by SopB | |
| | and ARHGEF26/SGEF, which induces cytoskeleton rearrangements and promotes bacterial | |
| | entry. | |
| Gene ID: | 391 | |
| UniProt: | P84095 | |
| Pathways: | RTK Signaling | |
| Application Details | | |
| Application Notes: | FCM 1:20-100 | |
| | IF(IHC-P) 1:50-200 | |
| | IF(IHC-F) 1:50-200 | |
| | IF(ICC) 1:50-200 | |
| Restrictions: | For Research Use only | |
| Handling | | |
| Format: | Liquid | |
| Concentration: | 1 μg/μL | |
| Buffer: | Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and | |
| | 50 % Glycerol. | |
| Preservative: | ProClin | |
| Precaution of Use: | This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be | |
| | handled by trained staff only. | |
| Storage: | -20 °C | |
| Storage Comment: | Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles. | |
| Expiry Date: | 12 months | |