# antibodies -online.com





## anti-ZBTB6 antibody (AbBy Fluor® 488)



|  | Go to Pi | oduct page |
|--|----------|------------|
|  |          |            |

| / N    | 11/0  | K /  | $\sim$ | A /         |
|--------|-------|------|--------|-------------|
|        | ve    | 1 \/ | ı⊢۱    | $\Lambda I$ |
| $\sim$ | ' V C | 1 V  |        | v v         |

| Quantity:    | 100 μL   |
|--------------|--|
| Target:      | ZBTB6  |
| Reactivity:  | Human, Mouse, Rat                                    |
| Host:        | Rabbit   |
| Clonality:   | Polyclonal   |
| Conjugate:   | This ZBTB6 antibody is conjugated to AbBy Fluor® 488 |
| Application: | Western Blotting (WB)                                |

#### **Product Details**

| Immunogen:            | KLH conjugated synthetic peptide derived from human ZBTB6/ZNF482 |
|-----------------------|--|
| Isotype:              | IgG  |
| Cross-Reactivity:     | Human  |
| Predicted Reactivity: | Mouse,Rat  |
| Purification:         | Purified by Protein A.   |

### Target Details

| Target:           | ZBTB6  |
|-------------------|--|
| Alternative Name: | Zbtb6/Znf482 (ZBTB6 Products)  |
| Background:       | Synonyms: ZBTB 6, ZBTB6, ZBTB6_HUMAN, ZID, Zinc finger and BTB domain containing                 |
|                   | protein 6, Zinc finger and BTB domain-containing protein 6, Zinc finger protein 482, Zinc finger |

protein with interaction domain, ZNF 482, ZNF482.

Background: Zinc-finger proteins contain DNA-binding domains and have a wide variety of functions, most of which encompass some form of transcriptional activation or repression. The majority of zinc-finger proteins contain a Kr\_ppel-type DNA binding domain and a KRAB domain, which is thought to interact with KAP1, thereby recruiting histone modifying proteins. Zinc finger and BTB domain-containing protein 5 (ZBTB5) is a 677 amino acid member of the Kr\_ppel C2H2-type zinc-finger protein family. Localized to the nucleus, ZBTB5 contains a BTB domain, also known as a POZ domain, which inhibits DNA binding and mediates homotypic and heterotypic dimerization. Characteristics of the BTB domain suggest that ZBTB5 functions as a transcription regulator.

Gene ID:

10773

#### **Application Details**

| Application Notes: | IF(IHC-P) 1:50-200    |
|--------------------|-----------------------|
| Restrictions:      | For Research Use only |
| Handling           |                       |
| Format:            | Liquid                |

| i oiiiidt.         | Liquid  |
|--------------------|---|
| Concentration:     | 1 μg/μL   |
| Buffer:            | Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and 50 % Glycerol.              |
| Preservative:      | Sodium azide  |
| Precaution of Use: | This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only. |
| Storage:           | -20 °C  |
| Storage Comment:   | Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.                                       |
| Expiry Date:       | 12 months   |