# antibodies -online.com





Datasheet for ABIN1695024

# anti-TESK2 antibody (AA 301-400) (Alexa Fluor 488)



( )	1/0	r\ /1	014	
( )	ve	I V I	-v	V

Quantity:	100 μL
Target:	TESK2
Binding Specificity:	AA 301-400
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This TESK2 antibody is conjugated to Alexa Fluor 488
Application:	Western Blotting (WB), Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence (Paraffin-embedded Sections) (IF (p))

#### **Product Details**

Immunogen:	KLH conjugated synthetic peptide derived from human TESK2
Isotype:	IgG
Predicted Reactivity:	Human,Mouse,Rat,Horse,Chicken,Rabbit
Purification:	Purified by Protein A.

## **Target Details**

Target:	TESK2
Alternative Name:	TESK2 (TESK2 Products)
Background:	Synonyms: Dual specicity testis specic protein kinase 2, Dual specicity testis-specic protein

kinase 2,	MGC29168, OTTH	IUMP000000009093	s, RP23-109A3.2,	Tesk2, TESK2	2_HUMAN,
Testicula	r protein kinase 2.	Testis specic kinas	se 2		

Background: TESK2 is a nuclear protein that belongs to the protein kinase superfamily and is expressed in testis and prostate. Functioning as a dual-specificity protein kinase, TESK2 catalyzes the ATP-dependent phosphorylation of substrates and autophosphorylation on tyrosine and serine/threonine residues, thereby mediating intracellular signal transduction pathways. TESK2 requires magnesium as a cofactor and its catalytic activity is thought to play an important role in meiotic events such as spermatogenesis. TESK2 contains one protein kinase domain that is 65 % identical to the kinase domain found in TESK1 (testicular protein kinase 1), suggesting a similar role for these proteins in phosphorylation events. Three isoforms of TESK2 are expressed due to alternative splicing.

Pathways: Cell-Cell Junction Organization

#### **Application Details**

Application Notes:	IF(IHC-P) 1:50-200
	IF(IHC-F) 1:50-200
	IF(ICC) 1:50-200

Restrictions: For Research Use only

### Handling

Format:	Liquid
Concentration:	1 μg/μL
Buffer:	Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and 50 % Glycerol.
Preservative:	ProClin
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only.
Storage:	-20 °C
Storage Comment:	Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.
Expiry Date:	12 months