antibodies -online.com





anti-GMEB2 antibody (AA 101-200) (Alexa Fluor 488)



\sim			
	N/P	r\/I	i⊢₩

Quantity:	100 μL	
Target:	GMEB2	
Binding Specificity:	AA 101-200	
Reactivity:	Mouse	
Host:	Rabbit	
Clonality:	Polyclonal	
Conjugate:	This GMEB2 antibody is conjugated to Alexa Fluor 488	
Application:	Western Blotting (WB), Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence (Paraffin-embedded Sections) (IF (p))	

Product Details

Immunogen:	KLH conjugated synthetic peptide derived from human GMEB2	
Isotype:	IgG	
Cross-Reactivity:	Mouse	
Predicted Reactivity:	Human,Rat,Dog,Cow,Sheep,Horse,Chicken,Rabbit	
Purification:	Purified by Protein A.	

Target Details

Target:	GMEB2
Alternative Name:	GMEB2 (GMEB2 Products)

Target Details

rarget betails		
Background:	Synonyms: GMEB2, DNA binding protein p79P, DNA-binding protein p79P, Glucocorticoid	
	modulatory element binding protein 2, Glucocorticoid modulatory element-binding protein 2,	
	GMEB 2, GMEB-2, Gmeb2, GMEB2_HUMAN, KIAA1269, P79P, Parvovirus initiation factor p79, P	
	p79, P79, RP4 663D7.1.	
	Background: GMEB-2 is a DNA-binding protein that plays a crucial role modulating transcription	
	upon activation by steroid hormones. GMEB-2 is ubiquitously expressed with preferential	
	expression in developmentally important tissues, such as testis, bone marrow, placenta and	
	fetal tissues. It localizes to the nucleus and cytoplasm and contains a SAND domain near its N-	
	terminus and a C-terminal coiled coil structure. GMEB-2 functions as a homodimer or as a	
	heterodimer with GMEB-1. The formed complex specifically binds to glucocorticoid modulatory	
	elements (GME) in the promoter region of target genes and recruits the histone acetylase CREB	
	binding protein (CBP) during glucocorticoid signaling. This acts to increase sensitivity to low	
	concentrations of glucocorticoids. In addition, GMEB-2 functions as an auxiliary factor required	
	for parvovirus replication.	
Gene ID:	26205	
Application Details		
Application Notes:	IF(IHC-P) 1:50-200	
	IF(IHC-F) 1:50-200	
	IF(ICC) 1:50-200	
Restrictions:	For Research Use only	
Handling		
Format:	Liquid	
Concentration:	1 μg/μL	
Buffer:	Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and 50 % Glycerol.	
Preservative:	ProClin	
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only.	

Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.

-20 °C

Storage:

Storage Comment:

\vdash	land	lına
	iaria	шц

Expiry Date:

12 months