# antibodies -online.com





## anti-CRTAP antibody (AA 201-300) (Alexa Fluor 488)



Go to Product page

( )	11/0	K\ /	iew	1
	$\cup$	'I V/I	$\square \vee \vee$	ı

Quantity:	100 μL
Target:	CRTAP
Binding Specificity:	AA 201-300
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This CRTAP antibody is conjugated to Alexa Fluor 488
Application:	Western Blotting (WB), Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence (Paraffin-embedded Sections) (IF (p))

#### **Product Details**

Immunogen:	KLH conjugated synthetic peptide derived from human CRTAP
Isotype:	IgG
Cross-Reactivity:	Human
Predicted Reactivity:	Mouse,Rat,Dog,Pig,Horse,Chicken,Rabbit
Purification:	Purified by Protein A.

#### **Target Details**

Target:	CRTAP
Alternative Name:	CRTAP (CRTAP Products)

### **Target Details**

Expiry Date:

ranget Betane			
Background:	Synonyms: Cartilage associated protein, LEPREL3, leprecan-like 3, CASP, CRTAP_HUMAN.		
	Background: CRTAP is a secreted protein localizing to the extracellular space that plays a role in		
	collagen post-translational modifications, extracellular fibril assembly and intracellular		
	trafficking. CRTAP is widely expressed with predominant expression in articular chondrocytes.		
	It contains a signal peptide and a tetratricopeptide-like helical domain and is essential for		
	normal bone formation. In the endoplasmic reticulum (ER), CRTAP forms a complex with Gros1		
	and CyPB (cyclophilin B) and is required for the efficient 3-hydroxylation of target prolyl residues		
	in Collagen Type I molecules, the major structural proteins of skin and bone. Mutations in the		
	gene encoding CRTAP can lead to autosomal recessive osteogenesis imperfecta (OI) type 7		
	and type 2B. OI, also known as brittle bone disease, is characterized by bone fragility and		
	susceptibility to fractures. OI type 7 is a mild form of this disorder, while OI type 2B is a neonatal		
	lethal condition.		
Gene ID:	10491		
Application Details			
Application Notes:	IF(IHC-P) 1:50-200		
	IF(IHC-F) 1:50-200		
	IF(ICC) 1:50-200		
Restrictions:	For Research Use only		
Handling			
Format:	Liquid		
Concentration:	1 μg/μL		
Buffer:	Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and		
	50 % Glycerol.		
Preservative:	ProClin		
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be		
	handled by trained staff only.		
Storage:	-20 °C		
Storage Comment:	Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.		

12 months