

## Datasheet for ABIN1695216

## anti-AKTIP antibody (AA 201-292) (AbBy Fluor® 488)



## Overview

| Overview              |  |
|-----------------------|--|
| Quantity:             | 100 μL   |
| Target:               | AKTIP  |
| Binding Specificity:  | AA 201-292   |
| Reactivity:           | Rat  |
| Host:                 | Rabbit   |
| Clonality:            | Polyclonal   |
| Conjugate:            | This AKTIP antibody is conjugated to AbBy Fluor® 488   |
| Application:          | Western Blotting (WB), Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence (Paraffin-embedded Sections) (IF (p)) |
| Product Details       |  |
| Immunogen:            | KLH conjugated synthetic peptide derived from human FTS/AKTIP  |
| Isotype:              | IgG  |
| Cross-Reactivity:     | Rat  |
| Predicted Reactivity: | Human,Mouse,Dog,Cow,Sheep,Pig,Horse,Chicken,Rabbit   |
| Purification:         | Purified by Protein A.   |
| Target Details        |  |
| Target:               | AKTIP  |
| Alternative Name:     | FTS/AKTIP (AKTIP Products)   |

## **Target Details**

| Background:        | Synonyms: AKT interacting protein, AKT-interacting protein, AKTIP, AKTIP_HUMAN, FT 1, FT1,   |  |
|--------------------|--|--|
|                    | Fused toes homolog, Fused toes protein homolog.  |  |
|                    | Background: Fused toes protein homolog (FTS), also known as AKT-interacting protein (AKTIP)  |  |
|                    | and Ft1, is a 292 amino acid protein that localizes to the cytoplasm and the cell membrane. A  |  |
|                    | member of the ubiquitin-conjugating enzyme family, FTS binds directly to AKT1 to regulate  |  |
|                    | apoptosis in a cell population. AKT1 is a protein that plays a critical role in a number of cellular   |  |
|                    | responses, such as cell growth, protein synthesis, and antiapoptotic signaling. The interaction of FTS and AKT1 enhances the phosphorylation and activation of AKT1, which, through an AKT1/GSK-3/NFATc1 signaling cascade, results in the increased production of the |  |
|                    |  | proapoptotic hormone Fas ligand and thus an increase in apoptosis. |
|                    |  | Gene ID:   |
|                    | Application Details  |  |
| Application Notes: | IF(IHC-P) 1:50-200   |  |
|                    | IF(IHC-F) 1:50-200   |  |
|                    | IF(ICC) 1:50-200   |  |
| Restrictions:      | For Research Use only  |  |
| Handling           |  |  |
| Format:            | Liquid   |  |
| Concentration:     | 1 μg/μL  |  |
| Buffer:            | Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and  |  |
|                    | 50 % Glycerol.   |  |
| Preservative:      | ProClin  |  |
| Precaution of Use: | This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be  |  |
|                    | handled by trained staff only.   |  |
| Storage:           | -20 °C   |  |
| Storage Comment:   | Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.  |  |
| Expiry Date:       | 12 months  |  |
|                    |  |  |