

Datasheet for ABIN1695244

anti-HIC2 antibody (AA 501-600) (AbBy Fluor® 488)[Go to Product page](#)

Overview

Quantity:	100 µL
Target:	HIC2
Binding Specificity:	AA 501-600
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This HIC2 antibody is conjugated to AbBy Fluor® 488
Application:	Western Blotting (WB), Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence (Paraffin-embedded Sections) (IF (p))

Product Details

Immunogen:	KLH conjugated synthetic peptide derived from human HIC2
Isotype:	IgG
Predicted Reactivity:	Human, Mouse, Rat, Dog, Cow, Pig, Horse, Chicken
Purification:	Purified by Protein A.

Target Details

Target:	HIC2
Alternative Name:	Hic2 (HIC2 Products)
Background:	Synonyms: HIC1 related gene on chromosome 22, HIC2, Hic3, HRG22, Hypermethylated in

Target Details

cancer 2 protein, KIAA1020, ZBTB30, Zinc finger and BTB domain-containing protein 30, ZNF907, HIC2_HUMAN.

Background: Zinc-finger proteins contain DNA-binding domains and have a wide variety of functions, most of which encompass some form of transcriptional activation or repression. The majority of zinc-finger proteins contain a Kr_ppl-type DNA binding domain and a KRAB domain, which is thought to interact with KAP1, thereby recruiting histone modifying proteins. HIC-2 (hypermethylated in cancer 2) possesses zinc finger motifs that are thought to be important for DNA-binding and also has a BTB/POZ domain at the N-terminus, which is thought to be important for protein-protein binding, as well as for the binding of transcription factors. HIC-2 is also known as Hic-3, HIC1-related gene on chromosome 22 or Zinc finger and BTB domain-containing protein 30, and is a 615 amino acid protein that is expressed as two isoforms produced by alternative splicing. HIC-2 is highly expressed in cerebellum and is localized to the nucleus in cells. HIC-2 contains a short amino acid sequence that is thought to interact with CtBP, a transcriptional repressor. The gene sequence associated with HIC-2 is thought to be a target for miRNAs (microRNAs) which are expressed in many cancers, suggesting that HIC-2 could possess tumor suppressor capabilities.

Gene ID: 23119

Application Details

Application Notes: IF(IHC-P) 1:50-200
IF(IHC-F) 1:50-200
IF(ICC) 1:50-200

Restrictions: For Research Use only

Handling

Format: Liquid

Concentration: 1 µg/µL

Buffer: Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and 50 % Glycerol.

Preservative: ProClin

Precaution of Use: This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only.

Handling

Storage:	-20 °C
Storage Comment:	Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.
Expiry Date:	12 months