

Datasheet for ABIN1695415

anti-FBLN2 antibody (AA 501-600) (AbBy Fluor® 488)[Go to Product page](#)

Overview

| | |
|----------------------|--|
| Quantity: | 100 µL |
| Target: | FBLN2 |
| Binding Specificity: | AA 501-600 |
| Reactivity: | Human |
| Host: | Rabbit |
| Clonality: | Polyclonal |
| Conjugate: | This FBLN2 antibody is conjugated to AbBy Fluor® 488 |
| Application: | Western Blotting (WB), Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence (Paraffin-embedded Sections) (IF (p)) |

Product Details

| | |
|-----------------------|---|
| Immunogen: | KLH conjugated synthetic peptide derived from human Fibulin 2 |
| Isotype: | IgG |
| Predicted Reactivity: | Human, Mouse, Rat, Cow, Sheep |
| Purification: | Purified by Protein A. |

Target Details

| | |
|-------------------|---|
| Target: | FBLN2 |
| Alternative Name: | Fibulin 2 (FBLN2 Products) |
| Background: | Synonyms: Fibulin-2, Fbln2, FBLN2_HUMAN, FIBL-2, Fibulin-2, OTTHUMP00000216670, |

Target Details

OTTHUMP00000216671, OTTHUMP00000216672.

Background: Fibulin-1 and Fibulin-2 associate with Fibronectin and other extracellular matrix proteins. In bone marrow, Fibulin-1 and Fibulin-2 bind to Fibronectin in the adherent layer. Fibulin-1 expression is stimulated by estrogen in ovarian cancer cell lines and has been suggested as both an agent of metastasis in ovarian cancer cells and as an indicator for predicting cancer risk or aggressiveness in ovarian carcinomas. The mobility of cancer cells may be inhibited with increasing exposure to Fibulin-1. Fibulin-2 binds to the lectin domains of extracellular matrix proteins aggrecan, versican and brevican. Fibulin-2 is abundantly expressed in heart, placenta and ovarian tissue, where it localizes to basement membranes and connective tissue compartments. In mice, differential Fibulin-2 gene expression correlates with the early phase of diabetic kidneys and glomerulosclerosis. The gene encoding human Fibulin-2 maps to chromosome 3p25.1.

Gene ID: 2199

Application Details

Application Notes: IF(IHC-P) 1:50-200
IF(IHC-F) 1:50-200
IF(ICC) 1:50-200

Restrictions: For Research Use only

Handling

Format: Liquid

Concentration: 1 µg/µL

Buffer: Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and 50 % Glycerol.

Preservative: ProClin

Precaution of Use: This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only.

Storage: -20 °C

Storage Comment: Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.

Expiry Date: 12 months