

Datasheet for ABIN1695457

anti-EIF2A antibody (pSer52) (AbBy Fluor® 488)



[Go to Product page](#)

Overview

Quantity:	100 µL
Target:	EIF2A
Binding Specificity:	pSer52
Reactivity:	Human, Mouse, Rat
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This EIF2A antibody is conjugated to AbBy Fluor® 488
Application:	Western Blotting (WB), Flow Cytometry (FACS), Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence (Paraffin-embedded Sections) (IF (p))

Product Details

Immunogen:	KLH conjugated synthetic phosphopeptide derived from human eIF2 alpha subunit 1 around the phosphorylation site of Ser52 [EL(p-S)RR]
Isotype:	IgG
Cross-Reactivity:	Human, Mouse, Rat
Predicted Reactivity:	Cow
Purification:	Purified by Protein A.

Target Details

Target:	EIF2A
---------	-------

Target Details

Alternative Name:	eIF2 alpha (EIF2A Products)
Background:	Synonyms: EIF2, EIF-2, EIF2A, EIF-2A, EIF-2alpha, Eukaryotic translation initiation factor 2 subunit 1, Eukaryotic translation initiation factor 2 subunit alpha, eIF-2-alpha, EIF2S1 Background: Functions in the early steps of protein synthesis by forming a ternary complex with GTP and initiator tRNA. This complex binds to a 4S ribosomal subunit, followed by mRNA binding to form a 43S preinitiation complex. Junction of the 6S ribosomal subunit to form the 8S initiation complex is preceded by hydrolysis of the GTP bound to eIF-2 and release of an eIF-2-GDP binary complex. In order for eIF-2 to recycle and catalyze another round of initiation, the GDP bound to eIF-2 must exchange with GTP by way of a reaction catalyzed by eIF-2B.
Gene ID:	1965
UniProt:	P05198
Pathways:	Ribonucleoprotein Complex Subunit Organization , ER-Nucleus Signaling , Hepatitis C , Methionine Biosynthetic Process , Ribosome Assembly

Application Details

Application Notes:	FCM 1:20-100 IF(IHC-P) 1:50-200 IF(IHC-F) 1:50-200 IF(ICC) 1:50-200
Restrictions:	For Research Use only

Handling

Format:	Liquid
Concentration:	1 µg/µL
Buffer:	Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and 50 % Glycerol.
Preservative:	ProClin
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only.
Storage:	-20 °C
Storage Comment:	Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.

Handling

Expiry Date: 12 months