## antibodies -online.com





## anti-ATPase Inhibitory Factor 1 antibody (AA 26-106) (AbBy Fluor® 488)



Go to Product page

$\sim$					
	۱۱/	r١	/1	0	۱۸/

Alternative Name:

Background:

Overview		
Quantity:	100 μL	
Target:	ATPase Inhibitory Factor 1 (ATPIF1)	
Binding Specificity:	AA 26-106	
Reactivity:	Rat	
Host:	Rabbit	
Clonality:	Polyclonal	
Conjugate:	This ATPase Inhibitory Factor 1 antibody is conjugated to AbBy Fluor® 488	
Application:	Western Blotting (WB)	
Product Details		
Immunogen:	KLH conjugated synthetic peptide derived from human ATPIF1/ATPase Inhibitory Factor 1	
Isotype:	IgG	
Cross-Reactivity:	Human	
Predicted Reactivity:	Mouse,Rat,Dog,Horse,Rabbit	
Purification:	Purified by Protein A.	
Target Details		
Target:	ATPase Inhibitory Factor 1 (ATPIF1)	

Synonyms: AT1\_HUMAN, ATP synthase inhibitor protein, ATPase inhibitor, ATPase inhibitor

ATP1/ATPase Inhibitory Factor 1 (ATPIF1 Products)

mitochondrial, ATPase inhibitor protein, ATPase inhibitory factor 1, ATPI, ATP1, ATP1, ATPIP, 1,
1, Inhibitor of F1Fo-ATPase, IP, MGC1167, MGC8898, mitochondrial.

Background: Mitochondrial ATP synthases (ATPases) transduce the energy contained in membrane electrochemical proton gradients into the energy required for synthesis of highenergy phosphate bonds. ATPases contain two linked complexes: F1, the hydrophilic catalytic core, and F0, the membrane-embedded protein channel. F1 consists of three Alpha chains and three Beta chains, which are weakly homologous, as well as one Gamma chain, one Delta chain and one Gamma chain. F0 consists of three subunits: a, b and c. A mitochondrial F1-ATPase inhibitor protein, ATPIF1 (ATPase inhibitory factor 1), also known as IP, IF1, ATPI or ATPIP (ATPase inhibitor protein), binds to the C-terminal region of a Beta subunit of the F1-ATPase at low pH values and, via interference of the Beta and Gamma subunit interaction, ATPIF1 regulates the activity of the F1F0-ATPase. This reversible ATPIF1 binding to F1F0-ATPase also occurs on the surface of endothelial cells.

Gene ID:

93974

## **Application Details**

Application Notes:	IF(IHC-P) 1:50-200
	15(1110 5) 4 50 000

IF(IHC-F) 1:50-200 IF(ICC) 1:50-200

Restrictions:

For Research Use only

## Handling

Format:	Liquid
Concentration:	1 μg/μL
Buffer:	Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and 50 % Glycerol.
Preservative:	ProClin
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only.
Storage:	-20 °C
Storage Comment:	Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.
Expiry Date:	12 months