



[Go to Product page](#)

Datasheet for ABIN1695543  
**anti-BTF3 antibody (AA 101-206) (AbBy Fluor® 488)**

### Overview

|                      |  |
|----------------------|--|
| Quantity:            | 100 µL   |
| Target:              | BTF3   |
| Binding Specificity: | AA 101-206   |
| Reactivity:          | Human  |
| Host:                | Rabbit   |
| Clonality:           | Polyclonal   |
| Conjugate:           | This BTF3 antibody is conjugated to AbBy Fluor® 488  |
| Application:         | Western Blotting (WB), Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence (Paraffin-embedded Sections) (IF (p)) |

### Product Details

|                       |  |
|-----------------------|--|
| Immunogen:            | KLH conjugated synthetic peptide derived from human BTF3 |
| Isotype:              | IgG  |
| Predicted Reactivity: | Human, Mouse, Rat, Dog, Cow, Sheep, Horse, Zebrafish     |
| Purification:         | Purified by Protein A.                                   |

### Target Details

|                   |  |
|-------------------|--|
| Target:           | BTF3   |
| Alternative Name: | BTF3 ( <a href="#">BTF3 Products</a> )   |
| Background:       | Synonyms: basic transcription factor 3, BETA NAC, BTF 3, btf3, BTF3_HUMAN, BTF3a, BTF3b, |

## Target Details

---

butyrophilin, subfamily 3, member A3, NACB, Nascent polypeptide associated complex beta polypeptide, RNA polymerase B transcription factor 3, Transcription factor BTF3.  
Background: General transcription factor. BTF3 can form a stable complex with RNA polymerase II. Required for the initiation of transcription.

## Application Details

---

Application Notes: IF(IHC-P) 1:50-200  
IF(IHC-F) 1:50-200  
IF(ICC) 1:50-200

Restrictions: For Research Use only

## Handling

---

Format: Liquid

Concentration: 1 µg/µL

Buffer: Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and 50 % Glycerol.

Preservative: ProClin

Precaution of Use: This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only.

Storage: -20 °C

Storage Comment: Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.

Expiry Date: 12 months