

[Go to Product page](#)

Datasheet for ABIN1695852

anti-GEM antibody (AA 201-296) (Alexa Fluor 488)

Overview

| | |
|----------------------|---|
| Quantity: | 100 µL |
| Target: | GEM |
| Binding Specificity: | AA 201-296 |
| Reactivity: | Human |
| Host: | Rabbit |
| Clonality: | Polyclonal |
| Conjugate: | This GEM antibody is conjugated to Alexa Fluor 488 |
| Application: | Western Blotting (WB), Flow Cytometry (FACS), Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence (Paraffin-embedded Sections) (IF (p)) |

Product Details

| | |
|-----------------------|---|
| Immunogen: | KLH conjugated synthetic peptide derived from human GEM |
| Isotype: | IgG |
| Cross-Reactivity: | Human |
| Predicted Reactivity: | Mouse,Rat,Dog,Cow,Pig,Horse |
| Purification: | Purified by Protein A. |

Target Details

| | |
|-------------------|--------------------------------------|
| Target: | GEM |
| Alternative Name: | GEM (GEM Products) |

Target Details

| | |
|-------------|--|
| Background: | <p>Synonyms: GTP binding mitogen induced T cell protein, GTP binding protein expressed in mitogen stimulated T cells, GTP binding protein GEM, GTP binding protein overexpressed in skeletal muscle, Kinase inducible Ras like protein, KIR, MGC26294, RAS like protein KIR, GEM_HUMAN.</p> <p>Background: Gem belongs to the Rad/Gem/Kir (RGK) subfamily of Ras-related GTPases, which lack typical C-terminal amino acid motifs for isoprenylation. Rad and Gem bind calmodulin in a Ca²⁺-dependent manner via this C-terminal extension, involving residues 278297 in human Rad. High intracellular Gem levels, which interact with intact microtubules and microfilaments, promote profound changes in cell morphology. Ectopic Gem expression is sufficient to stimulate cell flattening and neurite extension in N1E-115 and SH-SY5Y neuroblastoma cells, suggesting a role for Gem in cytoskeletal rearrangement and/or morphological differentiation of neurons. Gem was also observed in developing trigeminal nerve ganglia in 12.5 day mouse embryos, demonstrating that Gem expression is a property of normal ganglionic development. The interaction of Gem with beta-subunits regulates Ca²⁺ channel expression at the cell surface. The human Gem gene maps to chromosome ,8q22.1.</p> |
|-------------|--|

| | |
|----------|------|
| Gene ID: | 2669 |
|----------|------|

Application Details

| | |
|--------------------|--|
| Application Notes: | FCM 1:20-100 IF(IHC-P) 1:50-200 IF(IHC-F) 1:50-200 IF(ICC) 1:50-200 |
|--------------------|--|

| | |
|---------------|-----------------------|
| Restrictions: | For Research Use only |
|---------------|-----------------------|

Handling

| | |
|--------------------|--|
| Format: | Liquid |
| Concentration: | 1 µg/µL |
| Buffer: | Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and 50 % Glycerol. |
| Preservative: | ProClin |
| Precaution of Use: | This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only. |

Handling

| | |
|------------------|---|
| Storage: | -20 °C |
| Storage Comment: | Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles. |
| Expiry Date: | 12 months |