

Datasheet for ABIN1696060

anti-FAU antibody (AA 11-74) (AbBy Fluor® 488)



[Go to Product page](#)

Overview

Quantity:	100 µL
Target:	FAU
Binding Specificity:	AA 11-74
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This FAU antibody is conjugated to AbBy Fluor® 488
Application:	Western Blotting (WB), Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence (Paraffin-embedded Sections) (IF (p))

Product Details

Immunogen:	KLH conjugated synthetic peptide derived from human FAU
Isotype:	IgG
Predicted Reactivity:	Human, Mouse, Rat, Cow, Sheep, Pig, Horse
Purification:	Purified by Protein A.

Target Details

Target:	FAU
Alternative Name:	FAU/MNSF-beta (FAU Products)
Background:	Synonyms: 40S ribosomal protein S30, asr1, FAU, FAU encoded ubiquitin like protein, FAU1,

Target Details

FBR MuSV associated ubiquitously expressed, Finkel Biskis Reilly murine sarcoma virus FBR MuSV ubiquitously expressed fox derived, Finkel Biskis Reilly murine sarcoma virus FBR MuSV ubiquitously expressed, Finkel Biskis Reilly murine sarcoma virus ubiquitously expressed, FLJ22986, Fub1, Fubi, MNSFbeta, Monoclonal nonspecific suppressor factor beta, Ribosomal protein S30, RPS30, S30, UBIM_HUMAN, Ubiquitin like protein fubi and ribosomal protein S30, Ubiquitin like protein fubi, Ubiquitin like S30 fusion protein, Ubiquitin-like protein FUBI.

Background: The ubiquitin (Ub) pathway involves three sequential enzymatic steps that facilitate the conjugation of Ub and Ub-like molecules to specific protein substrates. The first step requires the ATP-dependent activation of the Ub C-terminus and the assembly of multi-Ub chains by the Ub-activating enzyme known as the E1 component. The Ub chain is then conjugated to the Ub-conjugating enzyme (E2) to generate an intermediate Ub-E2 complex. The Ub-ligase (E3) then catalyzes the transfer of Ub from E2 to the appropriate protein substrate. A wide range of enzymes facilitate in the proteolytic Ub pathway, including monoclonal nonspecific suppressor factor-beta (MNSF-beta), a subunit of MNSF, which is a lymphokine product of a murine T cell hybrid-oma that restricts the production of LPS-induced immunoglobulin secreting cells in an antigen-nonspecific manner. MNSF-beta is a ubiquitin-like fusion protein consisting of the ribosomal protein S30 and a protein that shares 36 % sequence identity with ubiquitin. This ubiquitin-like segment (Ubi-L) can be cleaved from MNSF-beta in the cytosol.

Gene ID: 2197

UniProt: [P62861](#)

Application Details

Application Notes: IF(IHC-P) 1:50-200
IF(IHC-F) 1:50-200
IF(ICC) 1:50-200

Restrictions: For Research Use only

Handling

Format: Liquid

Concentration: 1 µg/µL

Buffer: Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and 50 % Glycerol.

Handling

Preservative:	ProClin
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only.
Storage:	-20 °C
Storage Comment:	Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.
Expiry Date:	12 months