

[Go to Product page](#)

Datasheet for ABIN1696076

**anti-GGA3 antibody (AA 51-150) (Alexa Fluor 488)**

## Overview

Quantity:	100 µL
Target:	GGA3
Binding Specificity:	AA 51-150
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This GGA3 antibody is conjugated to Alexa Fluor 488
Application:	Western Blotting (WB), Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence (Paraffin-embedded Sections) (IF (p))

## Product Details

Immunogen:	KLH conjugated synthetic peptide derived from human GGA3
Isotype:	IgG
Predicted Reactivity:	Human, Mouse, Rat, Cow, Sheep, Pig, Horse, Rabbit
Purification:	Purified by Protein A.

## Target Details

Target:	GGA3
Alternative Name:	GGA3 ( <a href="#">GGA3 Products</a> )
Background:	Synonyms: ADP ribosylation factor binding protein 3, ADP ribosylation factor binding protein

## Target Details

GGA 3, ADP ribosylation factor binding protein GGA3, ADP-ribosylation factor-binding protein GGA3, ARF binding protein GGA 3, ARF binding protein GGA3, ARF-binding protein 3, gamma ear-containing, GGA 3, GGA3, GGA3\_HUMAN, Golgi associated gamma adaptin ear containing ARF binding protein 3, Golgi localized gamma ear containing ARF binding protein 3, Golgi-localized, KIAA0154.

Background: The GGA family of proteins (Golgi-localized, g-Adaptin ear-containing, ARF-binding proteins) are ubiquitous coat proteins that facilitate the trafficking of soluble proteins from the trans-Golgi network (TGN) to endosomes/lysosomes by means of interactions with TGN-sorting receptors, ARF (ADP-ribosylation factor), and clathrin. Members of the GGA family, GGA1, GGA2 (also known as VEAR) and GGA3, are multi-domain proteins that bind mannose 6-phosphate receptors (MPRs). GGAs have modular structures with an N-terminal VHS (VPS27, Hrs and STAM) domain followed by a GAT (GGA and Tom1) domain, a connecting hinge segment and a C-terminal GAE (g-Adaptin ear) domain. The amino-terminal VHS domains of GGAs form complexes with the cytoplasmic domains of sorting receptors by recognizing acidic-cluster di-leucine (ACLL) sequences. The human GGA3 gene maps to chromosome 17 and encodes a 723 amino acid protein that shares 46 % sequence identity with GGA1 and 38 % with GGA2.

Gene ID: 23163

## Application Details

Application Notes: IF(IHC-P) 1:50-200  
IF(IHC-F) 1:50-200  
IF(ICC) 1:50-200

Restrictions: For Research Use only

## Handling

Format: Liquid

Concentration: 1 µg/µL

Buffer: Aqueous buffered solution containing 0.01M TBS ( pH 7.4) with 1 % BSA, 0.03 % Proclin300 and 50 % Glycerol.

Preservative: ProClin

Precaution of Use: This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only.

Handling

---

Storage:	-20 °C
Storage Comment:	Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.
Expiry Date:	12 months