antibodies -online.com





anti-GNPDA1 antibody (AA 201-289) (Alexa Fluor 488)



| _ | | | | | |
|---|---|----|----|---|---|
| U | V | er | VI | е | W |

| Quantity: | 100 μL |
|----------------------|--|
| Target: | GNPDA1 |
| Binding Specificity: | AA 201-289 |
| Reactivity: | Mouse |
| Host: | Rabbit |
| Clonality: | Polyclonal |
| Conjugate: | This GNPDA1 antibody is conjugated to Alexa Fluor 488 |
| Application: | Western Blotting (WB), Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence (Paraffin-embedded Sections) (IF (p)) |

Product Details

| Immunogen: | KLH conjugated synthetic peptide derived from human GNPDA1 |
|-----------------------|--|
| Isotype: | IgG |
| Cross-Reactivity: | Mouse |
| Predicted Reactivity: | Human,Rat,Cow,Sheep,Pig |
| Purification: | Purified by Protein A. |

Target Details

| Target: | GNPDA1 |
|-------------------|--------------------------|
| Alternative Name: | GNPDA1 (GNPDA1 Products) |

Target Details

| Background: | Synonyms: EC 3.5.99.6, GlcN6P deaminase 1, Glucosamine 6 phosphate deaminase 1, |
|---------------------|---|
| | Glucosamine 6 phosphate isomerase 1, Glucosamine-6-phosphate deaminase 1, Glucosamine- |
| | 6-phosphate isomerase 1, GNPDA 1, GNPDA1, GNPI, GNPI1_HUMAN, HLN, KIAA0060, Oscillin, |
| | Putative glucosamine 6 phosphate isomerase. |
| | Background: During fertilization in mammals, the sperm activates the egg by causing an |
| | increase in the level of free cytoplasmic calcium concentration. This increased calcium |
| | concentration induces a characteristic series of oscillations that trigger egg activation and early |
| | embryo development. A hamster protein named oscillin is thought to be involved in this |
| | pathway. The enzyme glucosamine-6-phosphate isomerase (GNPI) or deaminase (GNPDA1) |
| | and the related protein GNPDA2 are the human homologs of hamster oscillin. GNPDA1 and |
| | GNPDA2 catalyze the conversion of GNP to fructose-6-phosphate and ammonia. Both proteins |
| | exist as homohexamers and are ubiquitously expressed with highest expression in testis, ovary |
| | and heart. Three isoforms of GNPDA2 are expressed due to alternative splicing events. |
| Gene ID: | 10007 |
| | |
| Application Details | |
| Application Notes: | IF(IHC-P) 1:50-200 |
| | IF(IHC-F) 1:50-200 |
| | IF(ICC) 1:50-200 |
| Restrictions: | For Research Use only |
| Handling | |
| Format: | Liquid |
| Concentration: | 1 μg/μL |
| | |
| Buffer: | Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and |
| | 50 % Glycerol. |
| Preservative: | ProClin |
| Precaution of Use: | This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be |
| | handled by trained staff only. |
| Storage: | -20 °C |
| Storage Comment: | Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles. |
| Expiry Date: | 12 months |