

[Go to Product page](#)

Datasheet for ABIN1696346

**anti-C4B antibody (AA 1454-1550) (Alexa Fluor 555)**

## Overview

Quantity:	100 µL
Target:	C4B (C4b)
Binding Specificity:	AA 1454-1550
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This C4B antibody is conjugated to Alexa Fluor 555
Application:	Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence (Paraffin-embedded Sections) (IF (p))

## Product Details

Immunogen:	KLH conjugated synthetic peptide derived from human Complement C4 gamma chain
Isotype:	IgG
Predicted Reactivity:	Human
Purification:	Purified by Protein A.

## Target Details

Target:	C4B (C4b)
Alternative Name:	Complement C4 gamma chain ( <a href="#">C4b Products</a> )
Background:	Synonyms: C4, RG, C4S, CO4, C4A2, C4A3, C4A4, C4A6, C4AD, CPAMD2, CH, C4F, C4B1, C4B2,

## Target Details

C4B3, C4B5, C4BD, C4B12, C4B\_2, CPAMD3, Complement C4-A, Acidic complement C4, C3 and PZP-like alpha-2-macroglobulin domain-containing protein 2, C4A

Background: Non-enzymatic component of C3 and C5 convertases and thus essential for the propagation of the classical complement pathway. Covalently binds to immunoglobulins and immune complexes and enhances the solubilization of immune aggregates and the clearance of IC through CR1 on erythrocytes. C4A isotype is responsible for effective binding to form amide bonds with immune aggregates or protein antigens, while C4B isotype catalyzes the transacylation of the thioester carbonyl group to form ester bonds with carbohydrate antigens. Derived from proteolytic degradation of complement C4, C4a anaphylatoxin is a mediator of local inflammatory process. It induces the contraction of smooth muscle, increases vascular permeability and causes histamine release from mast cells and basophilic leukocytes.

Gene ID:	720721
UniProt:	<a href="#">P0C0L4</a>
Pathways:	<a href="#">Complement System</a>

## Application Details

Application Notes:	IF(IHC-P) 1:50-200 IF(IHC-F) 1:50-200 IF(ICC) 1:50-200
Restrictions:	For Research Use only

## Handling

Format:	Liquid
Concentration:	1 µg/µL
Buffer:	Aqueous buffered solution containing 0.01M TBS ( pH 7.4) with 1 % BSA, 0.03 % Proclin300 and 50 % Glycerol.
Preservative:	ProClin
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only.
Storage:	-20 °C
Storage Comment:	Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.

Handling

---

Expiry Date: 12 months