

Datasheet for ABIN1696354

anti-TYROBP antibody (AA 11-100) (AbBy Fluor® 555)



and the contract of the contra

| Overvious | |
|-----------------------|---|
| Overview | |
| Quantity: | 100 μL |
| Target: | TYROBP |
| Binding Specificity: | AA 11-100 |
| Reactivity: | Human |
| Host: | Rabbit |
| Clonality: | Polyclonal |
| Conjugate: | This TYROBP antibody is conjugated to AbBy Fluor® 555 |
| Application: | Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence (Paraffin-embedded Sections) (IF (p)) |
| Product Details | |
| Immunogen: | KLH conjugated synthetic peptide derived from human DAP12 |
| Isotype: | IgG |
| Predicted Reactivity: | Human |
| Purification: | Purified by Protein A. |
| Target Details | |
| Target: | TYROBP |
| Alternative Name: | DAP12 (TYROBP Products) |
| Background: | Synonyms: DAP 12, DAP12, DNAX activation protein 12, DNAX-activation protein 12, KAR- |

associated protein, KARAP, Killer activating receptor associated protein, Killer-activating receptor-associated protein, PLOSL, TYOBP_HUMAN, TYRO protein tyrosine kinase binding protein, TYRO protein tyrosine kinase-binding protein, TYROBP.

Background: This gene encodes a transmembrane signaling polypeptide which contains an immunoreceptor tyrosine-based activation motif (ITAM) in its cytoplasmic domain. The encoded protein may associate with the killer-cell inhibitory receptor (KIR) family of membrane glycoproteins and may act as an activating signal transduction element. This protein may bind zeta-chain (TCR) associated protein kinase 70 kDa (ZAP-70) and spleen tyrosine kinase (SYK) and play a role in signal transduction, bone modeling, brain myelination, and inflammation. Mutations within this gene have been associated with polycystic lipomembranous osteodysplasia with sclerosing leukoencephalopathy (PLOSL), also known as Nasu-Hakola disease. Its putative receptor, triggering receptor expressed on myeloid cells 2 (TREM2), also causes PLOSL. Multiple alternative transcript variants encoding distinct isoforms have been identified for this gene. [provided by RefSeq, Mar 2010]

Application Details

Application Notes:

IF(IHC-P) 1:50-200

| Application Notes. | 11 (110 1) 1.00 200 |
|--------------------|---|
| | IF(IHC-F) 1:50-200 |
| | IF(ICC) 1:50-200 |
| Restrictions: | For Research Use only |
| Handling | |
| Format: | Liquid |
| Concentration: | 1 μg/μL |
| Buffer: | Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and |
| | 50 % Glycerol. |
| Preservative: | ProClin |
| Precaution of Use: | This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be |
| | handled by trained staff only. |
| Storage: | -20 °C |
| Storage Comment: | Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles. |
| Expiry Date: | 12 months |