

[Go to Product page](#)

Datasheet for ABIN1696443

anti-RBMY1F antibody (AA 54-160) (Alexa Fluor 555)

Overview

Quantity:	100 µL
Target:	RBMY1F
Binding Specificity:	AA 54-160
Reactivity:	Rat
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This RBMY1F antibody is conjugated to Alexa Fluor 555
Application:	Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence (Paraffin-embedded Sections) (IF (p))

Product Details

Immunogen:	KLH conjugated synthetic peptide derived from human RBMY1F
Isotype:	IgG
Cross-Reactivity:	Rat
Predicted Reactivity:	Human
Purification:	Purified by Protein A.

Target Details

Target:	RBMY1F
Alternative Name:	RBMY1F (RBMY1F Products)

Target Details

Background:	<p>Synonyms: YRRM2, RNA binding mot protein, Y chromosome, family 1 member F/J, RNA binding mot protein, Y linked, family 1, member F, Y chromosome RNA recognition mot 2, YRRM2, RBY1F_HUMAN.</p> <p>Background: The RBM (RNA-binding motif) gene family encodes proteins with an RNA binding motif. RBMY (RBM, Y chromosome) encodes a germ-cell specific nuclear protein involved in spermatogenesis. The RBM gene family, including RBMY1A, RBMY1B, RBMY1D, RBMY1E, RBMY1F, RBMY1H and RBMY1J, is comprised of at least 30 genes and pseudogenes, found on both arms of the Y chromosome. RBM X, an ancestral X chromosome homolog of the RBMY gene, encodes hnRNP G, which is widely expressed, whereas the RBMY gene evolved a male-specific function in spermatogenesis. Micro-deletions of the AZFb region of the Y chromosome, which contains a number of RBMY genes, usually result in severe consequences for spermatogenesis. RBM expression is localized to the nuclei of germ cells and RBM interacts with Tra2beta. Tra2beta is a ubiquitous activator of pre-mRNA splicing, but is most highly expressed in testis, suggesting a role for RBM in Tra2beta-dependent splicing in spermatocytes. The human RBMX gene maps to chromosome Xq26 and the RBMY gene family is found on all mammalian Y chromosomes.</p>
-------------	--

Gene ID:	159163
----------	--------

Application Details

Application Notes:	IF(IHC-P) 1:50-200 IF(IHC-F) 1:50-200 IF(ICC) 1:50-200
--------------------	--

Restrictions:	For Research Use only
---------------	-----------------------

Handling

Format:	Liquid
Concentration:	1 µg/µL
Buffer:	Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and 50 % Glycerol.
Preservative:	ProClin
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only.

Handling

Storage:	-20 °C
Storage Comment:	Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.
Expiry Date:	12 months