

Datasheet for ABIN1696458

anti-TRIM22 antibody (AA 231-330) (AbBy Fluor® 555)[Go to Product page](#)

Overview

Quantity:	100 µL
Target:	TRIM22
Binding Specificity:	AA 231-330
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This TRIM22 antibody is conjugated to AbBy Fluor® 555
Application:	Western Blotting (WB), Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence (Paraffin-embedded Sections) (IF (p))

Product Details

Immunogen:	KLH conjugated synthetic peptide derived from human Trim22
Isotype:	IgG
Cross-Reactivity:	Human
Purification:	Purified by Protein A.

Target Details

Target:	TRIM22
Alternative Name:	Trim22/Staf-50 (TRIM22 Products)
Background:	Synonyms: 50 kDa stimulated trans acting factor, RING finger protein 94, RNF94, Staf 50,

Target Details

STAF50, Stimulated trans acting factor 50 kDa, Stimulated Trans Acting Factor homolog of Mouse Rpt 1 gene, Trim22, Tripartite binding mot 22, Tripartite mot containing protein 22, TTRIM22RIM22, TRI22_HUMAN.

Background: The tripartite motif (TRIM) family of proteins are characterized by a conserved TRIM domain that includes a coiled-coil region, a B-box type zinc finger, one RING finger and three zinc-binding domains. Staf-50 (50 kDa-stimulated trans-acting factor), also known as TRIM22 (tripartite motif-containing 22), RNF94 or GPSTAF50, is a 498 amino acid cytoplasmic protein that belongs to the TRIM family and, characteristic of TRIM family members, contains one RING-type zinc finger, one B box-type zinc finger and one SPRY domain. Induced by IFN- α , and IFN- γ , Staf-50 is strongly expressed in ovary, spleen, thymus and peripheral blood leukocytes where it is thought to mediate the antiviral effects of IFN proteins. Additionally, Staf-50 is present in leukemic cells, suggesting a role in cancer formation and metastasis. Staf-50 exists as two alternatively spliced isoforms which are encoded by a gene that maps to human chromosome 11.

Gene ID: 10346

Application Details

Application Notes: IF(IHC-P) 1:50-200
IF(IHC-F) 1:50-200
IF(ICC) 1:50-200

Restrictions: For Research Use only

Handling

Format: Liquid

Concentration: 1 $\mu\text{g}/\mu\text{L}$

Buffer: Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and 50 % Glycerol.

Preservative: ProClin

Precaution of Use: This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only.

Storage: -20 °C

Storage Comment: Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.

Handling

Expiry Date: 12 months