## antibodies .- online.com





## anti-Acetyl-CoA Carboxylase beta antibody (pSer220) (Alexa Fluor 555)



Go to Product page

( )\	ve	K\ /	01	Α .
1 11	$/ \square$	I \/ I	$\square$	/\/

Immunofluorescence (Paraffin-embedded Sections) (IF (p)), Immunofluorescence (Cultured Cells) (IF (cc))		
me A		
_		

## **Target Details**

Storage Comment:

rarget betails	
Background:	Synonyms: Acetyl Coenzyme A Carboxylase beta phospho S220, Acetyl Coenzyme A
	Carboxylase beta phospho Ser2210 p-Acetyl Coenzyme A Carboxylase beta phospho S220,
	ACACB_HUMAN
	Background: Acetyl-CoA carboxylase (ACC) is a complex multifunctional enzyme system. ACC
	is a biotin-containing enzyme which catalyzes the carboxylation of acetyl-CoA to malonyl-CoA,
	the rate-limiting step in fatty acid synthesis. There are two ACC forms, alpha and beta, encoded
	by two different genes. ACC-alpha is highly enriched in lipogenic tissues. The enzyme is under
	long term control at the transcriptional and translational levels and under short term regulation
	by the phosphorylation/dephosphorylation of targeted serine residues and by allosteric
	transformation by citrate or palmitoyl-CoA. Multiple alternatively spliced transcript variants
	divergent in the 5' sequence and encoding distinct isoforms have been found for this gene.
	[provided by RefSeq, Jul 2008].
UniProt:	000763
Pathways:	AMPK Signaling, Ribonucleoside Biosynthetic Process
Application Details	
Application Notes:	IF(IHC-P) 1:50-200
	IF(IHC-F) 1:50-200
	IF(ICC) 1:50-200
Restrictions:	For Research Use only
Handling	
Format:	Liquid
Concentration:	1 μg/μL
Buffer:	Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and
	50 % Glycerol.
Preservative:	ProClin
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be
	handled by trained staff only.
Storage:	-20 °C

Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.

		1.	
$\vdash$	land	lın	
	ıaııu	1111	U

Expiry Date:

12 months