# antibodies -online.com







# anti-FASTKD3 antibody (AA 451-550) (Alexa Fluor 555)



$\sim$			
	N/P	r\/	i⊢₩

Quantity:	100 μL
Target:	FASTKD3
Binding Specificity:	AA 451-550
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This FASTKD3 antibody is conjugated to Alexa Fluor 555
Application:	Western Blotting (WB), Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence (Paraffin-embedded Sections) (IF (p))

#### **Product Details**

Immunogen:	KLH conjugated synthetic peptide derived from human FAKD3	
Isotype:	IgG	
Predicted Reactivity:	Human,Mouse,Pig,Horse,Rabbit	
Purification:	Purified by Protein A.	

## **Target Details**

Target:	FASTKD3
Alternative Name:	FAKD3 (FASTKD3 Products)
Background: Synonyms: FAKD3_HUMAN, FAST kinase domain containing protein 3, FAST kinase domain	

containing protein 3, FAST kinase domains 3, Fastkd3, FLJ23274, MGC142123, MGC5297. Background: FAKD3 is a member of a small family of Fas-activated serine/threonine kinase domain (FASTKD) containing proteins that share an amino terminal mitochondrial targeting domain and multiple carboxy terminal FAST domains as well as a putative RNA-binding RAP domain. The members of this family are ubiquitously expressed and are generally most abundant in mitochondria-enriched tissues such as heart, skeletal muscle and brown-adipose tissue. Some members of this protein family may play a role in apoptosis. The protein encoded by this gene interacts with components of the mitochondrial respiratory and translation networks. A pseudogene of this gene is also present on chromosome 5. Alternative splicing results in multiple transcript variants.

Gene ID:

79072

### **Application Details**

Application Notes: IF(IHC-P) 1:50-200

IF(IHC-F) 1:50-200 IF(ICC) 1:50-200

Restrictions:

For Research Use only

#### Handling

Format:	Liquid	
Concentration:	1 μg/μL	
Buffer:	Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and 50 % Glycerol.	
Preservative:	ProClin	
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only.	
Storage:	-20 °C	
Storage Comment:	Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.	
Expiry Date:	12 months	