antibodies .- online.com







anti-DAPL1 antibody (AA 1-100) (Alexa Fluor 555)



| () | ve | K\ / | | |
|-----|--------|------|----|----|
| | \cup | 1 V/ | Щ. | V۷ |
| | | | | |

| Quantity: | 100 μL | |
|----------------------|---|--|
| Target: | DAPL1 | |
| Binding Specificity: | AA 1-100 | |
| Reactivity: | Human | |
| Host: | Rabbit | |
| Clonality: | Polyclonal | |
| Conjugate: | This DAPL1 antibody is conjugated to Alexa Fluor 555 | |
| Application: | Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence (Paraffin-embedded Sections) (IF (p)) | |

Product Details

| Immunogen: | KLH conjugated synthetic peptide derived from human DAPL1 |
|-----------------------|---|
| Isotype: | IgG |
| Predicted Reactivity: | Human,Mouse,Rat,Horse |
| Purification: | Purified by Protein A. |

Target Details

| Target: | DAPL1 |
|-------------------|--|
| Alternative Name: | DAPL1 (DAPL1 Products) |
| Background: | Synonyms: dapL1, DAPL1_HUMAN, death associated protein-like 1, Death-associated protein- |

| | like 1, Early epithelial dferentiation-associated protein, EEDA, RP23-151F21.4. |
|---|---|
| | Background: In contrast to growth factors which promote cell proliferation, FAS ligand (FAS-L) |
| | and the tumor necrosis factors (TNFs) rapidly induce apoptosis. Cellular response to FAS-L and |
| | TNF is mediated by structurally related receptors containing a conserved cytoplasmic region |
| | called the death domain?. DAPL1 (Death-associated protein-like 1), also known as EEDA (Early |
| | epithelial differentiation-associated protein), is a 107 amino acid protein that is expressed in |
| | hair follicles and is thought to function in a similar manner to DAP-1, possibly participating in |
| | the early stages of epithelial differentiation and/or apoptosis. |
| | 92196 |
| | |
| | |
| _ | IF(IHC-P) 1:50-200 |
| | IF(IHC-F) 1:50-200 |
| | IF(ICC) 1:50-200 |
| | |
| | For Research Use only |
| | |
| | |
| | Liquid |

Handling

Restrictions:

Gene ID:

Application Details

Application Notes:

| Format: | Liquid | |
|--------------------|--|--|
| Concentration: | 1 μg/μL | |
| Buffer: | Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and 50 % Glycerol. | |
| Preservative: | ProClin | |
| Precaution of Use: | This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only. | |
| Storage: | -20 °C | |
| Storage Comment: | nment: Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles. | |
| Expiry Date: | 12 months | |