antibodies .- online.com





anti-FXYD6 antibody (AA 15-75) (Alexa Fluor 555)



O -	4 -	D	4	page
	$T \cap$	$Pr \cap C$	шет	nane
\cup	w	1 100	IUCL	Dauc

\sim	
()\/△	rview
\cup	1 410 44

Quantity:	100 μL	
Target:	FXYD6	
Binding Specificity:	AA 15-75	
Reactivity:	Human	
Host:	Rabbit	
Clonality:	Polyclonal	
Conjugate:	This FXYD6 antibody is conjugated to Alexa Fluor 555	
Application:	Western Blotting (WB), Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence (Paraffin-embedded Sections) (IF (p))	

Product Details

Immunogen:	KLH conjugated synthetic peptide derived from human FXYD6	
Isotype:	IgG	
Predicted Reactivity:	Human,Mouse,Rat	
Purification:	Purified by Protein A.	

Target Details

Target:	FXYD6
Alternative Name:	FXYD6 (FXYD6 Products)
Background:	Synonyms: FXYD domain containing ion transport regulator 6, FXYD domain containing ion

transport regulator 6 precursor, Phosphohippolin, UNQ521/PRO1056, FXYD6_HUMAN. Background: The mammalian FXYD family maintains Na+ and K+ gradients between the intracellular and extracellular milieus of cells in processes such as renal Na+-reabsorption, muscle contraction and neuronal excitability. FXYDs are single-span membrane proteins that share a 35 amino acid signature domain, beginning with the sequence PFXYD and containing seven invariant and six conserved amino acids. Members of the FXYD family include FXYD1 (PLM, phospholemman), FXYD2 (the g subunit of the Na+/K+-ATPase), FXYD3 (Mat8, mammary tumor protein), FXYD4 (CHIF) and FXYD5 (RIC). FXYD6 is expressed in various epithelial cells bordering the endolymph space and in the auditory neurons. FXYD6 co-localizes with Na+/K+-ATPase in the stria vascularis and can be co-immunoprecipitated with Na+/K+-ATPase. After expression, FXYD6 associates with Na+/K+-ATPase alpha1-beta1 and alpha1-beta2 isozymes, which are preferentially expressed in different regions of the inner ear and also with gastric and non-gastric H+/K+-ATPase.

Gene ID:

53826

Application Details

Application	Notes:

IF(IHC-P) 1:50-200

IF(IHC-F) 1:50-200

IF(ICC) 1:50-200

Restrictions:

For Research Use only

Handling

Format:	Liquid
Concentration:	1 μg/μL
Buffer:	Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and 50 % Glycerol.
Preservative:	ProClin
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only.
Storage:	-20 °C
Storage Comment:	Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.
Expiry Date:	12 months