



[Go to Product page](#)

Datasheet for ABIN1696778  
**anti-SOCS5 antibody (AA 151-250) (Alexa Fluor 555)**

### Overview

Quantity:	100 µL
Target:	SOCS5
Binding Specificity:	AA 151-250
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This SOCS5 antibody is conjugated to Alexa Fluor 555
Application:	Western Blotting (WB), Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence (Paraffin-embedded Sections) (IF (p))

### Product Details

Immunogen:	KLH conjugated synthetic peptide derived from human SOCS5
Isotype:	IgG
Predicted Reactivity:	Human,Mouse,Rat,Cow,Sheep,Chicken
Purification:	Purified by Protein A.

### Target Details

Target:	SOCS5
Alternative Name:	SOCS5 ( <a href="#">SOCS5 Products</a> )
Background:	Synonyms: CIS-6, CIS6, CISH5, CISH6, Cytokine inducible SH2 containing protein 5, Cytokine

## Target Details

---

inducible SH2 protein 6, KIAA0671, SOCS-5, Suppressor of cytokine signaling 5.

Background: The eight members of the recently identified Suppressor of Cytokines Signaling (SOCS) family are SOCS1, SOCS2, SOCS3, SOCS4, SOCS5, SOCS6, SOCS7, and CIS. Structurally the SOCS proteins are composed of an N-terminal region of variable length and amino acid composition, a central SH2 domain, and a C-terminal motif called the SOCS box. The SOCS proteins appear to form part of a classical negative feedback loop that regulates cytokine signal transduction. Transcription of each of the SOCS genes occurs rapidly in vitro and in vivo in response to cytokines, and once produced, the various members of the SOCS family appear to inhibit signaling in different ways. During Th1 differentiation a reduction in the association of Jak1 with the IL4 Receptor correlated with the appearance of SOCS5. SOCS5 protein was preferentially expressed in committed Th1 cells and interacted with the cytoplasmic region of the IL4 Receptor alpha chain irrespective of receptor tyrosine phosphorylation. This unconventional interaction of SOCS5 protein with IL4 Receptor resulted in the inhibition of IL4-mediated signal transducer and activator of transcription-6 activation. T cells from transgenic mice constitutively expressing SOCS5 exhibited a significant reduction of IL4-mediated Th2 development. Therefore, the induced SOCS5 protein in Th1 differentiation environment may play an important role by regulating Th1 and Th2 balance.

Pathways: [JAK-STAT Signaling](#), [EGFR Signaling Pathway](#), [Positive Regulation of Immune Effector Process](#)

## Application Details

---

Application Notes: IF(IHC-P) 1:50-200  
IF(IHC-F) 1:50-200  
IF(ICC) 1:50-200

Restrictions: For Research Use only

## Handling

---

Format: Liquid

Concentration: 1 µg/µL

Buffer: Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and 50 % Glycerol.

Preservative: ProClin

Precaution of Use: This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only.

## Handling

---

Storage: -20 °C

---

Storage Comment: Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.

---

Expiry Date: 12 months