antibodies.com

Datasheet for ABIN1696875 anti-TUBGCP3 antibody (AA 401-500) (Alexa Fluor 555)



Overview

Quantity:	100 µL	
Target:	TUBGCP3	
Binding Specificity:	AA 401-500	
Reactivity:	Xenopus laevis	
Host:	Rabbit	
Clonality:	Polyclonal	
Conjugate:	This TUBGCP3 antibody is conjugated to Alexa Fluor 555	
Application:	Western Blotting (WB), Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence (Paraffin-embedded Sections) (IF (p))	

Product Details

Immunogen:	KLH conjugated synthetic peptide derived from human GCP3/Gamma-tubulin complex component 3
Isotype:	lgG
Cross-Reactivity:	Xenopus laevis
Predicted Reactivity:	Human,Mouse,Rat,Cow,Sheep,Chicken
Purification:	Purified by Protein A.
Target Details	

Target:

TUBGCP3

Order at www.antibodies-online.com | www.antikoerper-online.de | www.anticorps-enligne.fr | www.antibodies-online.cn International: +49 (0)241 95 163 153 | USA & Canada: +1 877 302 8632 | support@antibodies-online.com Page 1/3 | Product datasheet for ABIN1696875 | 03/08/2024 | Copyright antibodies-online. All rights reserved.

Target Details		
Alternative Name:	GCP3/Gamma-tubulin complex component 3 (TUBGCP3 Products)	
Background:	Synonyms: Gamma-ring complex protein 104 kDa, Gamma-tubulin complex component 3, GCP	
	3, GCP3, GCP3_HUMAN, h104p, hGCP3, hGrip104, hSpc98, SPBC98, Spc98p, Spindle pole body	
	protein Spc98 homolog, TUBGCP3, Tubulin gamma complex associated protein 3.	
	Background: The Gamma-Tubulin complex is composed of Gamma Tubulin and the Gamma-	
	Tubulin complex-associated proteins GCP2, GCP3, GCP4, GCP5 and GCP6, all of which are	
	essential components of microtubule organizing centers. Gamma-Tubulin complex	
	components are localized to both the centrosome, where they are involved in microtubule	
	nucleation, and to the cytoplasm, where they exist as soluble complexes that can be recruited	
	to the centrosome as needed. Although the GCP proteins are related, they have distinct roles	
	which contribute to the proper function of the Gamma-Tubulin complex. GCP3 (Gamma-	
	Tubulin complex component 3), also known as TUBGCP3 or SPBC98, localizes to the	
	centrosome and is a ubiquitously expressed 907 amino acid member of the Gamma-Tubulin	
	complex. Like GCP2 and Gamma Tubulin, GCP3 is conserved in all eukaryotes, suggesting that	
	it is part of a core unit involved in eukaryotic microtubule nucleation. Three isoforms of GCP3	
	exist due to alternative splicing events.	
Gene ID:	10426	
Pathways:	M Phase, SARS-CoV-2 Protein Interactome	
Application Details		
Application Notes:	IF(IHC-P) 1:50-200	
	IF(IHC-F) 1:50-200	
	IF(ICC) 1:50-200	
Restrictions:	For Research Use only	
Handling		
Format:	Liquid	
Concentration:	1 µg/µL	
Buffer:	Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and	
	50 % Glycerol.	
Preservative:	ProClin	
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be	

Order at www.antibodies-online.com | www.antikoerper-online.de | www.anticorps-enligne.fr | www.antibodies-online.cn International: +49 (0)241 95 163 153 | USA & Canada: +1 877 302 8632 | support@antibodies-online.com Page 2/3 | Product datasheet for ABIN1696875 | 03/08/2024 | Copyright antibodies-online. All rights reserved.

1.1		
	lond	lina
	land	

	handled by trained staff only.	
Storage:	-20 °C	
Storage Comment:	Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.	
Expiry Date:	12 months	