antibodies .- online.com





anti-FAM20C antibody (AA 351-450) (Alexa Fluor 555)



\sim							
	1//	$rac{1}{2}$	`\/	L	Р	1	٨

Quantity:	100 μL
Target:	FAM20C
Binding Specificity:	AA 351-450
Reactivity:	Rat
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This FAM20C antibody is conjugated to Alexa Fluor 555
Application:	Western Blotting (WB), Immunofluorescence (Paraffin-embedded Sections) (IF (p)), Immunofluorescence (Cultured Cells) (IF (cc))

Product Details

Immunogen:	KLH conjugated synthetic peptide derived from human FAM20C
Isotype:	IgG
Cross-Reactivity:	Rat
Predicted Reactivity:	Human,Mouse,Dog,Cow,Sheep,Horse,Chicken,Monkey
Purification:	Purified by Protein A.

Target Details

Target:	FAM20C
Alternative Name:	FAM20C (FAM20C Products)

Target Details

Storage Comment:

rarget Details	
Background:	Synonyms: C76981, Dentin matrix protein 4, DKFZp547C074, DMP-4, DMP4, DMP4_HUMAN,
	Extracellular serine/threonine protein kinase Fam20C, Fam20C, Family with sequence similarity
	20 member C, GEF CK, Golgi enriched fraction casein kinase, Protein FAM20C, RNS antibod.
	Background: The FAM20 proteins are a family of secreted proteins that regulate differentiation
	and function of hematopoietic and other tissues. FAM20C, also known as DMP4 (Dentin matrix
	protein 4), is a 570 amino acid secreted protein that binds calcium and may play a role in dentin
	mineralization. Defects in the gene encoding FAM20C are the cause of Raine syndrome (Lethal
	osteosclerotic bone dysplasia), an autosomal recessive osteosclerotic bone dysplasia, that is
	characterized by generalized osteosclerosis, microencephaly and craniofacial dysplasia.
	Usually, affected individuals survive only days or weeks. The mutations of the FAM20C gene
	include four nonsynomous base changes and four splice-site changes that have a detrimental
	affect on splicing.
Gene ID:	56975
UniProt:	Q8IXL6
Application Details	
Application Notes:	IF(IHC-P) 1:50-200
	IF(IHC-F) 1:50-200
	IF(ICC) 1:50-200
Restrictions:	For Research Use only
Handling	
Format:	Liquid
Concentration:	1 μg/μL
Buffer:	Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and
	50 % Glycerol.
Preservative:	ProClin
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be
	handled by trained staff only.
Storage:	-20 °C

Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.

		1.	
\vdash	land	lın	
	ıaııu	1111	U

Expiry Date:

12 months