antibodies .- online.com





Datasheet for ABIN1696970

anti-PISD antibody (Alexa Fluor 555)



()	1/0	r\ /1	014	
()	ve	I V I	-v	V

- OVERVIEW	
Quantity:	100 μL
Target:	PISD
Reactivity:	Human, Mouse, Rat
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This PISD antibody is conjugated to Alexa Fluor 555
Application:	Western Blotting (WB), Immunofluorescence (Paraffin-embedded Sections) (IF (p))
Product Details	

Immunogen:	KLH conjugated synthetic peptide derived from human PISD
Isotype:	IgG
Cross-Reactivity:	Human, Mouse, Rat
Purification:	Purified by Protein A.

Target Details

Target:	PISD
Alternative Name:	PISD (PISD Products)
Background:	Synonyms: DJ858B16, dJ858B16.2, DKFZp566G2246, Phosphatidylserine decarboxylase, Phosphatidylserine decarboxylase beta chain, Phosphatidylserine decarboxylase proenzyme, PISD, PISD_HUMAN, PSD, PSDC, PSSC.
	Background: Enzymes known as phosphatidylserine decarboxylases (PSDs) catalyze the

Target Details

formation of phosphatidylethanolamine from phosphatidylserine via phosphatidylserine decarboxylation. Type I PSDs contain LGST motifs and are found in bacteria and eukaryotic mitochondria, whereas type II PSDs contain GGST motifs and are found in eukaryotic endomembrane systems. PISD (phosphatidylserine decarboxylase), also known as phosphatidylserine decarboxylase proenzyme, PSDC, PSD, PSSC, DJ858B16, dJ858B16.2 or DKFZp566G2246, is a 408 amino acid a type I phosphatidylserine decarboxylase that localizes to the inner mitochondrial membrane. PISD contains a conserved LGST motif which is cleaved to produce two isoforms known as PISD and PISD _. PISD is capable of forming a heterodimer and is highly expressed in liver and testis. The gene encoding PISD maps to human chromosome 22q12.2.

Gene ID:

23761

IF(IHC-P) 1:50-200

Application Details

Application Notes:

Application Notes.	IF(InC-P) 1.50-200	
Restrictions:	For Research Use only	
Handling		
Format:	Liquid	
Concentration:	1 μg/μL	
Buffer:	Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and 50 % Glycerol.	
Preservative:	ProClin	
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only.	
Storage:	-20 °C	
Storage Comment:	Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.	
Expiry Date:	12 months	