antibodies -online.com





anti-GARNL1 antibody (AA 631-730) (Alexa Fluor 555)



Go to Product page

| \sim | | | |
|--------|-----|-----|-----|
| | N/P | r\/ | i⊢₩ |

| Quantity: | 100 μL |
|----------------------|---|
| Target: | GARNL1 (RALGAPA1) |
| Binding Specificity: | AA 631-730 |
| Reactivity: | Rat |
| Host: | Rabbit |
| Clonality: | Polyclonal |
| Conjugate: | This GARNL1 antibody is conjugated to Alexa Fluor 555 |
| Application: | Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence (Paraffin-embedded Sections) (IF (p)) |

Product Details

| Immunogen: | KLH conjugated synthetic peptide derived from human GARNL1 |
|-----------------------|--|
| Isotype: | IgG |
| Cross-Reactivity: | Rat |
| Predicted Reactivity: | Human, Mouse, Dog, Cow, Sheep, Pig, Horse, Chicken |
| Purification: | Purified by Protein A. |

Target Details

| Target: | GARNL1 (RALGAPA1) |
|-------------------|----------------------------|
| Alternative Name: | GARNL1 (RALGAPA1 Products) |

Target Details

| Background: | Synonyms: DKFZp566D133, DKFZp667F074, GAP related interacting partner to E12, GARN L1, | |
|----------------------|--|--|
| | GARNL 1, GRIPE, GTPase activating Rap/RanGAP domain like 1, KIAA0884, Tuberin like protein | |
| | 1, TULIP 1. TULIP1, RGPA1_HUMAN. | |
| | Background: GARNL1 is expressed during embryogenesis with E12. During development, | |
| | GARNL1 expression decreases, persisting at high levels only in neurons of the adult brain. | |
| | GARNL1 localizes to the cytoplasm where it may play a role regulating GTP hydrolysis of | |
| | proteins such as Ran and Rap. GARNL1 is imported to the nucleus via dimerization with E12. | |
| | GARNL1 interacts with the HLH region of E12 and may function to negatively regulate the | |
| | transcription of E12-dependent downstream target genes. This suggests that at least a portion | |
| | of the function of GARNL1 is dependent upon its association with E12. GARNL1 may also | |
| | associate with other HLH proteins and influence a variety of HLH signaling cascades. In adult | |
| | brain, GARNL1 activity does not involve E12 and therefore it may serve a different function in | |
| | developed neural tissue. | |
| Gene ID: | 253959 | |
| Application Details | | |
| Application Notes: | IF(IHC-P) 1:50-200 | |
| Application (Votes). | IF(IHC-F) 1:50-200 | |
| | IF(ICC) 1:50-200 | |
| | | |
| Restrictions: | For Research Use only | |
| Handling | | |
| Format: | Liquid | |
| Concentration: | 1 μg/μL | |
| Buffer: | Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and | |
| | 50 % Glycerol. | |
| Preservative: | ProClin | |
| Precaution of Use: | This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be | |
| | handled by trained staff only. | |
| Storage: | -20 °C | |
| Storage Comment: | Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles. | |
| Expiry Date: | 12 months | |
| | | |