

[Go to Product page](#)

Datasheet for ABIN1697154

anti-EIF1 antibody (AA 41-113) (Alexa Fluor 555)

Overview

Quantity:	100 µL
Target:	EIF1
Binding Specificity:	AA 41-113
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This EIF1 antibody is conjugated to Alexa Fluor 555
Application:	Western Blotting (WB), Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence (Paraffin-embedded Sections) (IF (p))

Product Details

Immunogen:	KLH conjugated synthetic peptide derived from human eIF1
Isotype:	IgG
Predicted Reactivity:	Human, Mouse, Rat, Pig, Horse, Chicken
Purification:	Purified by Protein A.

Target Details

Target:	EIF1
Alternative Name:	eIF1 (EIF1 Products)
Background:	Synonyms: A121, e1, E1_HUMAN, E1A, Eukaryotic translation initiation factor 1, ISO1, Protein

Target Details

translation factor SUI1 homolog, SUI1, Sui1iso1.

Background: In mammalian cells, translation is controlled at the level of polypeptide chain initiation by initiation factors. Eukaryotic translation initiation factor 1 (eIF1) is crucial for the scanning process in vitro. During the scanning process, eIF1 is a component of a complex involved in recognition of the initiator codon. Translation is also initiated by the role of eIF1 in regulating the activity of ribosomal subunits 43S, 48S and 40S. eIF1 enables 43S ribosomal complexes to discern between cognate and near-cognate initiation codons, sensing the nucleotide content of initiation codons. It is also a promotor, along with eukaryotic translation initiation factor 1A (eIF1A), for assembly of 48S ribosomal complexes at the initiation codon of a conventional capped mRNA. In addition, eIF1 and eIF1A, together with eukaryotic translation initiation factor 5 (eIF5), function in the formation of stable 40S ribosomal preinitiation complexes.

Gene ID: 10209

Application Details

Application Notes: IF(IHC-P) 1:50-200
IF(IHC-F) 1:50-200
IF(ICC) 1:50-200

Restrictions: For Research Use only

Handling

Format: Liquid

Concentration: 1 µg/µL

Buffer: Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and 50 % Glycerol.

Preservative: ProClin

Precaution of Use: This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only.

Storage: -20 °C

Storage Comment: Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.

Expiry Date: 12 months