

[Go to Product page](#)

Datasheet for ABIN1697162

**anti-ANAPC10 antibody (AA 31-130) (Alexa Fluor 555)**

## Overview

Quantity:	100 µL
Target:	ANAPC10
Binding Specificity:	AA 31-130
Reactivity:	Rat
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This ANAPC10 antibody is conjugated to Alexa Fluor 555
Application:	Western Blotting (WB), Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence (Paraffin-embedded Sections) (IF (p))

## Product Details

Immunogen:	KLH conjugated synthetic peptide derived from human APC10
Isotype:	IgG
Cross-Reactivity:	Rat
Predicted Reactivity:	Human, Mouse, Cow, Sheep, Pig, Chicken, Rabbit
Purification:	Purified by Protein A.

## Target Details

Target:	ANAPC10
Alternative Name:	APC10 ( <a href="#">ANAPC10 Products</a> )

## Target Details

Background:	<p>Synonyms: ANAPC 10, anapc10, Anaphase promoting complex 10, Anaphase promoting complex subunit 10, Anaphase-promoting complex subunit 10, Apc 10, APC10, APC10_HUMAN, Cyclosome subunit 10, DKFZP564L0562, Doc 1, Doc1, OTTHUMP00000220297, OTTHUMP00000220298, OTTHUMP00000220299, OTTHUMP00000220300, OTTHUMP00000220303.</p> <p>Background: Composed of more than ten subunits, the anaphase-promoting complex (APC) acts in a cell-cycle dependent manner to promote the separation of sister chromatids during the transition between metaphase and anaphase in mitosis. APC, or cyclosome, accomplishes this progression through the ubiquitination of mitotic cyclins and other regulatory proteins that are targeted for destruction during cell division. APC is phosphorylated, and thus activated, by protein kinases Cdk1/cyclin B and polo-like kinase (Plk). APC is under tight control by a number of regulatory factors, including CDC20, CDH1 and MAD2. Specifically, CDC20 and CDH1 directly bind to and activate the cyclin-ubiquitination activity of APCs. In contrast, MAD2 inhibits APC by forming a ternary complex with CDC20 and APC, thus preventing APC activation. APC10 contains a Doc1 homology domain, which is a beta-sandwich structure common to many other putative E3 ubiquitin ligases. APC10 binds to core APC subunits throughout the cell cycle. Specifically, APC10 binds to the C-terminus of CDC27/APC3. During mitosis, APC10 is localized in centrosomes and mitotic spindles. APC10 also localizes to kinetochores from prophase to anaphase, and to the midbody in telophase and cytokinesis.</p>
-------------	--

Gene ID:	10393
----------	-------

Pathways:	<a href="#">M Phase</a>
-----------	-------------------------

## Application Details

Application Notes:	IF(IHC-P) 1:50-200 IF(IHC-F) 1:50-200 IF(ICC) 1:50-200
--------------------	--

Restrictions:	For Research Use only
---------------	-----------------------

## Handling

Format:	Liquid
---------	--------

Concentration:	1 µg/µL
----------------	---------

Buffer:	Aqueous buffered solution containing 0.01M TBS ( pH 7.4) with 1 % BSA, 0.03 % Proclin300 and 50 % Glycerol.
---------	---

## Handling

---

Preservative:	ProClin
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only.
Storage:	-20 °C
Storage Comment:	Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.
Expiry Date:	12 months