

Datasheet for ABIN1697244

anti-FZD8 antibody (AbBy Fluor® 555)[Go to Product page](#)

Overview

Quantity:	100 µL
Target:	FZD8
Reactivity:	Human, Mouse, Rat
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This FZD8 antibody is conjugated to AbBy Fluor® 555
Application:	Western Blotting (WB), Immunofluorescence (Paraffin-embedded Sections) (IF (p))

Product Details

Immunogen:	KLH conjugated synthetic peptide derived from human Frizzled 8
Isotype:	IgG
Cross-Reactivity:	Human, Mouse, Rat
Purification:	Purified by Protein A.

Target Details

Target:	FZD8
Alternative Name:	Frizzled 8 (FZD8 Products)
Background:	Synonyms: Frizzled 8 seven transmembrane spanning receptor, frizzled 8, seven transmembrane spanning receptor, Frizzled family receptor 8, frizzled homolog 8 Drosophila, Frizzled homolog 8, Frizzled-8, FZ 8, Fz-8, FZ8, FZD 8, FZD8, FZD8_HUMAN, hFZ 8, hFZ8, HGNC4046, Homolog of Drosophila Frizzled 8.

Target Details

Background: The frizzled gene, originally identified in *Drosophila melanogaster*, is involved in the development of tissue polarity. The mammalian homolog of frizzled, as well as several secreted mammalian frizzled-related proteins (FRPs), have been described. The frizzled proteins contain seven transmembrane domains, a cysteine-rich domain in the extracellular region and a carboxy-terminal Ser/Thr-xxx-Val motif. They function as receptors for Wnt and are generally coupled to G proteins. The cysteine-rich domain of frizzled-8 blocks endogenous Wnts and the effects of Wnt-1 and Wnt-5 on proliferation. The mouse frizzled-8 gene, which encodes a Wnt receptor, is a potent cancer-associated activator of the Beta-catenin-TCF pathway. The frizzled-8 gene contains no introns. Frizzled-8 mRNA has been detected in fetal brain and kidney, and also in adult pancreas, skeletal muscle, kidney and heart. Frizzled is highly expressed in HeLa S3 (cervical uterus cancer) cells and A549 lung cancer cells.

Gene ID: 8325

Pathways: [WNT Signaling](#)

Application Details

Application Notes: FCM: (1:20-100), IF(IHC-P): (1:50-200)
Optimal working dilution should be determined by the investigator.

Restrictions: For Research Use only

Handling

Format:	Liquid
Concentration:	1 µg/µL
Buffer:	Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and 50 % Glycerol.
Preservative:	Sodium azide
Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Storage:	-20 °C
Storage Comment:	Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.
Expiry Date:	12 months