# antibodies .- online.com





## anti-GLI4 antibody (AA 271-376) (AbBy Fluor® 555)



Go to Product page

$\sim$					
	1/6	⊃r	\/I	$\triangle$	٨/

Quantity:	100 μL
Target:	GLI4
Binding Specificity:	AA 271-376
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This GLI4 antibody is conjugated to AbBy Fluor® 555
Application:	Western Blotting (WB), Immunofluorescence (Paraffin-embedded Sections) (IF (p)), Immunofluorescence (Cultured Cells) (IF (cc))

## **Product Details**

Immunogen:	KLH conjugated synthetic peptide derived from human GLI4
Isotype:	IgG
Cross-Reactivity:	Human
Predicted Reactivity:	Mouse,Rat,Cow,Pig,Horse
Purification:	Purified by Protein A.

## **Target Details**

Target:	GLI4
Alternative Name:	GLI4 (GLI4 Products)

## Target Details

Racko	round:
Dacke	n Our iu.

Synonyms: GLI 4, GLI Kruppel family member GLI 4, GLI Kruppel family member GLI4, HKR4, HKR4, HKR4, Protein, Krueppel related zinc finger protein 4, Krueppel-related zinc finger protein 4, Neural specic DNA binding protein, Oncogene HKR 4, Oncogene HKR4, Protein HKR 4, Protein HKR4, Zinc finger protein GLI 4, Zinc finger protein GLI4.

Background: Zinc-finger proteins contain DNA-binding domains and have a wide variety of functions, most of which encompass some form of transcriptional activation or repression. The majority of zinc-finger proteins contain a krueppel-type DNA binding domain and a KRAB domain, which is thought to interact with KAP1, thereby recruiting histone modifying proteins.

GLI-4, also known as HKR4, is a 376 amino acid protein that localizes to the nucleus and contains seven C2H2-type zinc fingers. Belonging to the krueppel C2H2-type zinc-finger protein family, GLI-4 may function as a transcriptional regulator, effectively activating or repressing the transcription of target genes. The gene encoding GLI-4 maps to human chromosome 8, which consists of nearly 146 million base pairs, houses more than 800 genes and is associated with a variety of diseases and malignancies.

Gene ID:

2738

#### **Application Details**

lication	

IF(IHC-P) 1:50-200

IF(IHC-F) 1:50-200

IF(ICC) 1:50-200

Restrictions:

For Research Use only

## Handling

Format:	Liquid
Concentration:	1 μg/μL
Buffer:	Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and 50 % Glycerol.
Preservative:	ProClin
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only.
Storage:	-20 °C
Storage Comment:	Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.

ш	land	linc	٩
	ıanıu	11111	J

Expiry Date:

12 months