# antibodies -online.com





#### Datasheet for ABIN1697916

## anti-KIF13B antibody (AA 351-450) (AbBy Fluor® 555)



( )	11/0	K\ /	iew
	$\cup$	'I V/I	$I \cap VV$

Quantity:	100 μL
Target:	KIF13B
Binding Specificity:	AA 351-450
Reactivity:	Mouse, Rat
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This KIF13B antibody is conjugated to AbBy Fluor® 555
Application:	Western Blotting (WB), Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence (Paraffin-embedded Sections) (IF (p))

### **Product Details**

Immunogen:	KLH conjugated synthetic peptide derived from human KIF13B
Isotype:	IgG
Cross-Reactivity:	Mouse, Rat
Predicted Reactivity:	Human,Cow,Sheep,Pig,Horse
Purification:	Purified by Protein A.

### **Target Details**

Target:	KIF13B
Alternative Name:	KIF13B (KIF13B Products)

#### **Target Details**

Bac	kar	ัดเ	ın	ď	
Duo		$\sim$	<i>a</i>	<u> </u>	۰

Synonyms: GAKIN, Guanylate kinase associated kinesin, KI13B\_HUMAN, KIAA0639, K 13B, K13B, Kinesin 13B, Kinesin family member 13B, Kinesin like protein GAKIN, Kinesin like protein K13B, Kinesin-like protein GAKIN, Kinesin-like protein K13B, OTTHUMP00000225649.

Background: KIF13B is also known as Kinesin-like protein GAKIN or GAKIN and is a 1,826 amino acid protein that is widely expressed in tissues, with highest expression in brain and kidney.

KIF13B is localized to the cytoplasm, as well as to the cytoskeleton, and is thought to be a microtubule-dependent motor protein which is able to bind to a variety of proteins in order to traffic them to various locations throughout the cell. KIF13B belongs to the kinesin-like protein family and possesses three domains typical of the kinesin-like protein family, namely an N-terminal motor domain with an ATP-binding motif, an FHA domain which is known to bind diverse cargos and a large stalk domain involved in protein-protein binding. Additionally, KIF13B has a microtubule-interacting sequence which is known as the CAP-Gly domain at its C-terminus. The CAP-Gly domain is highly conserved domain among eukaryotes, and in humans, defects in the CAP-Gly domain are implicated in many diseases affecting the trafficking of vesicles, neuromuscular junctions and lysosome proliferation.

Gene ID:	23303	
UniProt:	Q9NQT8	

#### **Application Details**

<b>Application</b>	Notes:

IF(IHC-P) 1:50-200

IF(IHC-F) 1:50-200

IF(ICC) 1:50-200

Restrictions:

For Research Use only

#### Handling

Format:	Liquid
Concentration:	1 μg/μL
Buffer:	Aqueous buffered solution containing 0.01M TBS ( pH 7.4) with 1 % BSA, 0.03 % Proclin300 and 50 % Glycerol.
Preservative:	ProClin
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only.

## Handling

Storage:	-20 °C
Storage Comment:	Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.
Expiry Date:	12 months