antibodies .- online.com





anti-C10RF50 antibody (AA 111-199) (Alexa Fluor 555)



Go to Product page

| \sim | | | | | |
|--------|----|----|----|-------------|---|
| () | VE | ۲۱ | /1 | \triangle | Λ |

| Quantity: | 100 μL |
|----------------------|--|
| Target: | C10RF50 |
| Binding Specificity: | AA 111-199 |
| Reactivity: | Human |
| Host: | Rabbit |
| Clonality: | Polyclonal |
| Conjugate: | This C10RF50 antibody is conjugated to Alexa Fluor 555 |
| Application: | Western Blotting (WB), Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence (Paraffin-embedded Sections) (IF (p)) |

Product Details

| Immunogen: | KLH conjugated synthetic peptide derived from human C1orf50 |
|-----------------------|---|
| Isotype: | IgG |
| Predicted Reactivity: | Human,Rat,Rabbit |
| Purification: | Purified by Protein A. |

Target Details

| Target: | C10RF50 |
|-------------------|---|
| Alternative Name: | C1orf50 (C1ORF50 Products) |
| Background: | Synonyms: C1orf50, CA050_HUMAN, Chromosome 1 open reading frame 50, MGC955, |

Uncharacterized protein C1orf50.

Background: Chromosome 1 is the largest human chromosome spanning about 260 million base pairs and making up 8 % of the human genome. There are about 3,000 genes on chromosome 1, and considering the great number of genes there are also a large number of diseases associated with chromosome 1. Notably, the rare aging disease Hutchinson-Gilford progeria is associated with the LMNA gene which encodes lamin A. When defective, the LMNA gene product can build up in the nucleus and cause characteristic nuclear blebs. The mechanism of rapidly enhanced aging is unclear and is a topic of continuing exploration. The MUTYH gene is located on chromosome 1 and is partially responsible for familial adenomatous polyposis. Stickler syndrome, Parkinsons, Gaucher disease and Usher syndrome are also associated with chromosome 1. A breakpoint has been identified in 1q which disrupts the DISC1 gene and is linked to schizophrenia. Aberrations in chromosome 1 are found in a variety of cancers including head and neck cancer, malignant melanoma and multiple myeloma. The C1orf50 gene product has been provisionally designated C1orf50 pending further characterization.

| Gene ID: | 79078 |
|----------|-------|
| 5 | |

Pathways: SARS-CoV-2 Protein Interactome

Application Details

| Application Notes: | IF(IHC-P) 1:50-200 |
|--------------------|--------------------|
| | IF(IHC-F) 1:50-200 |
| | IF(ICC) 1:50-200 |
| | |

Restrictions: For Research Use only

Handling

| Format: | Liquid |
|--------------------|--|
| Concentration: | 1 μg/μL |
| Buffer: | Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and 50 % Glycerol. |
| Preservative: | ProClin |
| Precaution of Use: | This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only. |

Handling

| Storage: | -20 °C |
|------------------|---|
| Storage Comment: | Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles. |
| Expiry Date: | 12 months |