

Datasheet for ABIN1698102

**anti-Complement C4 antibody (AA 21-120) (AbBy Fluor® 647)**[Go to Product page](#)

## Overview

|                      |                                                                                                         |
|----------------------|---------------------------------------------------------------------------------------------------------|
| Quantity:            | 100 µL                                                                                                  |
| Target:              | Complement C4 (C4)                                                                                      |
| Binding Specificity: | AA 21-120                                                                                               |
| Reactivity:          | Human                                                                                                   |
| Host:                | Rabbit                                                                                                  |
| Clonality:           | Polyclonal                                                                                              |
| Conjugate:           | This Complement C4 antibody is conjugated to AbBy Fluor® 647                                            |
| Application:         | Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence (Paraffin-embedded Sections) (IF (p)) |

## Product Details

|                       |                                                                   |
|-----------------------|-------------------------------------------------------------------|
| Immunogen:            | KLH conjugated synthetic peptide derived from human Complement C4 |
| Isotype:              | IgG                                                               |
| Predicted Reactivity: | Human                                                             |
| Purification:         | Purified by Protein A.                                            |

## Target Details

|                   |                                                                                        |
|-------------------|----------------------------------------------------------------------------------------|
| Target:           | Complement C4 (C4)                                                                     |
| Alternative Name: | C4a+C4b ( <a href="#">C4 Products</a> )                                                |
| Background:       | Synonyms: C4, RG, C4S, CO4, C4A2, C4A3, C4A4, C4A6, C4AD, CPAMD2, CH, C4F, C4B1, C4B2, |

## Target Details

C4B3, C4B5, C4BD, C4B12, C4B\_2, CPAMD3, Complement C4-A, Acidic complement C4, C3 and PZP-like alpha-2-macroglobulin domain-containing protein 2, C4A

Background: Non-enzymatic component of C3 and C5 convertases and thus essential for the propagation of the classical complement pathway. Covalently binds to immunoglobulins and immune complexes and enhances the solubilization of immune aggregates and the clearance of IC through CR1 on erythrocytes. C4A isotype is responsible for effective binding to form amide bonds with immune aggregates or protein antigens, while C4B isotype catalyzes the transacylation of the thioester carbonyl group to form ester bonds with carbohydrate antigens. Derived from proteolytic degradation of complement C4, C4a anaphylatoxin is a mediator of local inflammatory process. It induces the contraction of smooth muscle, increases vascular permeability and causes histamine release from mast cells and basophilic leukocytes.

|           |                                   |
|-----------|-----------------------------------|
| Gene ID:  | 720, 721                          |
| UniProt:  | <a href="#">P0C0L4</a>            |
| Pathways: | <a href="#">Complement System</a> |

## Application Details

|                    |                                                              |
|--------------------|--------------------------------------------------------------|
| Application Notes: | IF(IHC-P) 1:50-200<br>IF(IHC-F) 1:50-200<br>IF(ICC) 1:50-200 |
| Restrictions:      | For Research Use only                                        |

## Handling

|                    |                                                                                                                    |
|--------------------|--------------------------------------------------------------------------------------------------------------------|
| Format:            | Liquid                                                                                                             |
| Concentration:     | 1 µg/µL                                                                                                            |
| Buffer:            | Aqueous buffered solution containing 0.01M TBS ( pH 7.4) with 1 % BSA, 0.03 % Proclin300 and 50 % Glycerol.        |
| Preservative:      | ProClin                                                                                                            |
| Precaution of Use: | This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only. |
| Storage:           | -20 °C                                                                                                             |
| Storage Comment:   | Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.                                  |

## Handling

---

Expiry Date: 12 months