# antibodies .- online.com





## anti-FANCM antibody (AA 831-930) (Alexa Fluor 647)



Go to Product page

| $\sim$ |     |     |     |
|--------|-----|-----|-----|
|        | N/P | r\/ | i⊢₩ |

| Quantity:            | 100 μL  |  |
|----------------------|---|--|
| Target:              | FANCM   |  |
| Binding Specificity: | AA 831-930  |  |
| Reactivity:          | Human   |  |
| Host:                | Rabbit  |  |
| Clonality:           | Polyclonal  |  |
| Conjugate:           | This FANCM antibody is conjugated to Alexa Fluor 647  |  |
| Application:         | Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence (Paraffin-embedded Sections) (IF (p)) |  |

#### **Product Details**

| Immunogen:            | KLH conjugated synthetic peptide derived from human FANCM |  |
|-----------------------|---|--|
| Isotype:              | IgG   |  |
| Predicted Reactivity: | Human   |  |
| Purification:         | Purified by Protein A.                                    |  |

### **Target Details**

| Target:           | FANCM  |  |
|-------------------|--|--|
| Alternative Name: | FANCM (FANCM Products)   |  |
| Background:       | Synonyms: FAAP250, Fanconi anemia group M protein, Protein Hef ortholog, |  |

Background: Fanconi anemia (FA) is an autosomal recessive disorder characterized by bone marrow failure, birth defects and chromosomal instability. At the cellular level, FA is characterized by spontaneous chromosomal breakage and a unique hypersensitivity to DNA cross-linking agents. The thirteen FA proteins that have been characterized are important for regulating chromosomal stability and genome surveillance. Eight of these proteins, namely FANCA, FANCB, FANCC, FANCE, FANCF, FANCG, FANCL and FANCM, comprise the FA core complex, which catalyzes a key reaction in DNA repair: the monoubiquitination of FANCD2. FANCM (Fanconi anemia, complementation group M) is a member of the DEAD-box helicase family of proteins and contains a DEAH helicase domain and a nuclease domain. Localizing to chromatin fractions, FANCM is phosphorylated in a cell cycle-dependent manner and is believed to function as an anchor, recruiting the FA core complex to chromatin. Mutations in the gene encoding FANCM can lead to Fanconi anemia.

Gene ID: 57697

Pathways: DNA Damage Repair

#### **Application Details**

Application Notes: IF(IHC-P) 1:50-200

IF(IHC-F) 1:50-200

IF(ICC) 1:50-200

Restrictions: For Research Use only

#### Handling

| Format:            | Liquid   |
|--------------------|--|
| Concentration:     | 1 μg/μL  |
| Buffer:            | Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and 50 % Glycerol.         |
| Preservative:      | ProClin  |
| Precaution of Use: | This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only. |
| Storage:           | -20 °C   |
| Storage Comment:   | Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.                                  |

| $\vdash$ | land  | lına |
|----------|-------|------|
|          | iaria | шц   |

Expiry Date:

12 months