antibodies .- online.com





anti-FUT6 antibody (AA 101-200) (AbBy Fluor® 647)



C ~ .	+ ~	1500	110+	page
(¬()	I() F	$T \cap T$	11 16 11	112016

()	1/0	r\ /1	014	
()	ve	I V I	-v	V

Quantity:	100 μL
Target:	FUT6
Binding Specificity:	AA 101-200
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This FUT6 antibody is conjugated to AbBy Fluor® 647
Application:	Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence (Paraffin-embedded Sections) (IF (p))

Product Details

Immunogen:	KLH conjugated synthetic peptide derived from human FUT6/FucT-VI
Isotype:	IgG
Predicted Reactivity:	Human
Purification:	Purified by Protein A.

Target Details

Target:	FUT6
Alternative Name:	FUT6 (FUT6 Products)
Background:	Synonyms: Alpha 1,3 fucosyltransferase, EC=2.4.1.65, FCT3A, Fuc TVI, Fucosyltransferase 6,

Fucosyltransferase VI, FucT VI, Galactoside 3 L fucosyltransferase, FUT6_HUMAN. Background: Glycosyltransferases that mediate the regio- and stereoselective transfer of sugars, such as the fucosyltransferases, determine cell surface-carbohydrate profiles, which is an essential interface for biological recognition processes. Fucosyltransferases catalyze the covalent association of fucose to different positional linkages in sugar acceptor molecules. The carbohydrate moieties generated and covalently attached to cell surfaces are necessary to ensure a surface contour that satisfies physiological roles, which are reliant on adhesion molecules such as Selectins (1-3). Hematopoietic lineages rely on Fucosyltransferases to confer a surface carbohydrate phenotype, which mediates proper cell adhesion molecule recruitment and cell trafficking (4-6).

Gene ID:

2528

Application Details

Application Notes: IF(IHC-P) 1:50-200
IF(IHC-F) 1:50-200

IF(ICC) 1:50-200

Restrictions:

For Research Use only

Handling

Format:	Liquid	
Concentration:	1 μg/μL	
Buffer:	Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and 50 % Glycerol.	
Preservative:	ProClin	
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only.	
Storage:	-20 °C	
Storage Comment:	Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.	
Expiry Date:	12 months	