# antibodies -online.com





# anti-TDRP/C8orf42 antibody (AA 51-150) (Alexa Fluor 647)



Go to Product page

$\sim$			
	N/P	r\/	i⊢₩

Quantity:	100 μL	
Target:	TDRP/C8orf42 (TDRP)	
Binding Specificity:	AA 51-150	
Reactivity:	Rat	
Host:	Rabbit	
Clonality:	Polyclonal	
Conjugate:	This TDRP/C8orf42 antibody is conjugated to Alexa Fluor 647	
Application:	Western Blotting (WB), Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence (Paraffin-embedded Sections) (IF (p))	

## **Product Details**

Immunogen:	KLH conjugated synthetic peptide derived from human C8orf42
Isotype:	IgG
Cross-Reactivity:	Rat
Predicted Reactivity:	Human,Mouse
Purification:	Purified by Protein A.

## **Target Details**

Target:	TDRP/C8orf42 (TDRP)
Alternative Name:	C8orf42 (TDRP Products)

## **Target Details**

Synonyms: C8orf42, TDRP_HUMAN, Protein INM01, Protein TDRP, Uncharacterized protein C8orf42.	
Background: Made up of nearly 146 million bases, chromosome 8 encodes about 800 genes.	
Translocation of portions of chromosome 8 with amplifications of the c-Myc gene are found in	
some leukemias and lymphomas, and typically associated with a poor prognosis. Portions of	
chromosome 8 have been linked to schizophrenia and bipolar disorder. Trisomy 8, also known	
as Warkany syndrome 2, most often results in early miscarriage but is occasionally seen in a	
mosaic form in surviving patients who suffer to a varying degree from a number of symptoms	
including retarded mental and motor development, and certain facial and developmental	
defects. WRN is a DNA helicase encoded by chromosome 8 and shown defective in those with	
the early aging disorder Werner syndrome. Chromosome 8 is also associated with Pfeiffer	
syndrome, congenital hypothyroidism and Waardenburg syndrome. The C8orf42 gene product	
has been provisionally designated C8orf42 pending further characterization.	
157695	
IF(IHC-P) 1:50-200	
IF(IHC-F) 1:50-200	
IF(ICC) 1:50-200	
For Research Use only	
Liquid	
1 μg/μL	
Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and	
50 % Glycerol.	
ProClin	
This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be	
handled by trained staff only.	
-20 °C	
Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.	
12 months	