

Datasheet for ABIN1698662

anti-TESK2 antibody (AA 301-400) (AbBy Fluor® 647)



Overview

| Overview | | |
|-----------------------|--|--|
| Quantity: | 100 μL | |
| Target: | TESK2 | |
| Binding Specificity: | AA 301-400 | |
| Reactivity: | Human | |
| Host: | Rabbit | |
| Clonality: | Polyclonal | |
| Conjugate: | This TESK2 antibody is conjugated to AbBy Fluor® 647 | |
| Application: | Western Blotting (WB), Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence (Paraffin-embedded Sections) (IF (p)) | |
| Product Details | | |
| Immunogen: | KLH conjugated synthetic peptide derived from human TESK2 | |
| Isotype: | IgG | |
| Predicted Reactivity: | Human,Mouse,Rat,Horse,Chicken,Rabbit | |
| Purification: | Purified by Protein A. | |
| Target Details | | |
| Target: | TESK2 | |
| Alternative Name: | TESK2 (TESK2 Products) | |
| Background: | Synonyms: Dual specicity testis specic protein kinase 2, Dual specicity testis-specic protein | |
| | | |

| kinase 2, MGC29168, OTTHUMP00000009093, RP23-109A3.2, Tesk2, TESK2_HUMAN, | |
|---|--|
| Testicular protein kinase 2, Testis specic kinase 2. | |

Background: TESK2 is a nuclear protein that belongs to the protein kinase superfamily and is expressed in testis and prostate. Functioning as a dual-specificity protein kinase, TESK2 catalyzes the ATP-dependent phosphorylation of substrates and autophosphorylation on tyrosine and serine/threonine residues, thereby mediating intracellular signal transduction pathways. TESK2 requires magnesium as a cofactor and its catalytic activity is thought to play an important role in meiotic events such as spermatogenesis. TESK2 contains one protein kinase domain that is 65 % identical to the kinase domain found in TESK1 (testicular protein kinase 1), suggesting a similar role for these proteins in phosphorylation events. Three isoforms of TESK2 are expressed due to alternative splicing.

Pathways: Cell-Cell Junction Organization

Application Details

| Application Notes: | IF(IHC-P) 1:50-200 | |
|--------------------|--------------------|--|
| | IF(IHC-F) 1:50-200 | |
| | IF(ICC) 1:50-200 | |

Restrictions: For Research Use only

Handling

| Format: | Liquid |
|--------------------|--|
| Concentration: | 1 μg/μL |
| Buffer: | Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and 50 % Glycerol. |
| Preservative: | ProClin |
| Precaution of Use: | This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only. |
| Storage: | -20 °C |
| Storage Comment: | Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles. |
| Expiry Date: | 12 months |