antibodies -online.com





Datasheet for ABIN1698666

anti-NGEF antibody (pTyr179) (Alexa Fluor 647)

Go to	Proc	luct	ра	ae

Overview	
Quantity:	100 μL
Target:	NGEF
Binding Specificity:	pTyr179
Reactivity:	Human, Mouse, Rat
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This NGEF antibody is conjugated to Alexa Fluor 647
Application:	Immunofluorescence (Paraffin-embedded Sections) (IF (p)), Immunofluorescence (Cultured Cells) (IF (cc)), Western Blotting (WB)
Product Details	
Immunogen:	KLH conjugated synthetic phosphopeptide derived from human Ephexin-1 around the phosphorylation site of Tyr179
Isotype:	IgG
Cross-Reactivity:	Human, Mouse, Rat
Predicted Reactivity:	Chicken
Purification:	Purified by Protein A.
Target Details	
Target:	NGEF

Target Details

nyms: Ephexin-1 phospho Y179, p-Ephexin-1 phospho Y179, ARHGEF27, Eph-interacting ange protein, EPHEXIN, Ephexin 1, Neuronal guanine nucleotide exchange factor, Ngef, F_HUMAN. ground: EPHEXIN is a 710 amino acid protein that localizes to both the membrane and the plasm and contains one SH3 domain, one PH domain and one DH domain. Expressed at levels in brain and present at lower levels in lung tissue, EPHEXIN interacts with EphA4 and cions as a guanine nucleotide exchange factor (GEF) that is capable of activating Rho A, I and Cdc42 and is thought to play a role in axon guidance and growth cone collapse. EXIN is subject to Src-dependent phosphorylation, an event that increases the GEF activity PHEXIN toward Rho A. Human EPHEXIN, which exists as multiple alternatively spliced rms, shares a high degree of sequence homology with its mouse counterpart, suggesting
plasm and contains one SH3 domain, one PH domain and one DH domain. Expressed at levels in brain and present at lower levels in lung tissue, EPHEXIN interacts with EphA4 and tions as a guanine nucleotide exchange factor (GEF) that is capable of activating Rho A, and Cdc42 and is thought to play a role in axon guidance and growth cone collapse. EXIN is subject to Src-dependent phosphorylation, an event that increases the GEF activity PHEXIN toward Rho A. Human EPHEXIN, which exists as multiple alternatively spliced
levels in brain and present at lower levels in lung tissue, EPHEXIN interacts with EphA4 and tions as a guanine nucleotide exchange factor (GEF) that is capable of activating Rho A, and Cdc42 and is thought to play a role in axon guidance and growth cone collapse. EXIN is subject to Src-dependent phosphorylation, an event that increases the GEF activity PHEXIN toward Rho A. Human EPHEXIN, which exists as multiple alternatively spliced
tions as a guanine nucleotide exchange factor (GEF) that is capable of activating Rho A, and Cdc42 and is thought to play a role in axon guidance and growth cone collapse. EXIN is subject to Src-dependent phosphorylation, an event that increases the GEF activity PHEXIN toward Rho A. Human EPHEXIN, which exists as multiple alternatively spliced
and Cdc42 and is thought to play a role in axon guidance and growth cone collapse. EXIN is subject to Src-dependent phosphorylation, an event that increases the GEF activity PHEXIN toward Rho A. Human EPHEXIN, which exists as multiple alternatively spliced
EXIN is subject to Src-dependent phosphorylation, an event that increases the GEF activity PHEXIN toward Rho A. Human EPHEXIN, which exists as multiple alternatively spliced
PHEXIN toward Rho A. Human EPHEXIN, which exists as multiple alternatively spliced
rms, shares a high degree of sequence homology with its mouse counterpart, suggesting
nserved role between species.
1
otrophin Signaling Pathway
C-P) 1:50-200
C-F) 1:50-200
C) 1:50-200
Research Use only
d
/μL
ous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and
Glycerol.
lin
product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be led by trained staff only.
 C

Handling

Storage Comment:	Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.
Expiry Date:	12 months