antibodies - online.com







anti-GNG5 antibody (AA 21-65) (Alexa Fluor 647)



_	
()Ver	view

Quantity:	100 μL
Target:	GNG5
Binding Specificity:	AA 21-65
Reactivity:	Rat
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This GNG5 antibody is conjugated to Alexa Fluor 647
Application:	Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence (Paraffin-embedded Sections) (IF (p))

Product Details

Immunogen:	KLH conjugated synthetic peptide derived from human GNG5
Isotype:	IgG
Cross-Reactivity:	Rat
Predicted Reactivity:	Human,Mouse,Cow,Chicken,Rabbit
Purification:	Purified by Protein A.

Target Details

Target:	GNG5
Alternative Name:	GNG5 (GNG5 Products)

Target Details

Background:	Synonyms: FLJ92393, GBG5_HUMAN, Gng5, Guanine nucleotide-binding protein GI/GS/GO
	subunit gamma-5, OTTHUMP00000011474, OTTHUMP00000011565.
	Background: Heterotrimeric G proteins function to relay information from cell surface receptors
	to intracellular effectors. Each of a very broad range of receptors specifically detects an
	extracellular stimulus (i.e. a photon, pheromone, odorant, hormone or neurotransmitter), while
	the effectors (e.g. adenyl cyclase), which act to generate one or more intracellular messengers,
	are less numerous. In mammals, G protein alhfa, beta and gamma polypeptides are encoded by
	at least 16, 4 and 7 genes, respectively. Most interest in G proteins has been focused on their a
	subunits, since these proteins bind and hydrolyze GTP and most obviously regulate the activity
	of the best studied effectors. Evidence, however, has established an important regulatory role
	for the beta gamma subunits. It is becoming increasingly clear that different G protein
	complexes expressed in different tissues carry structurally distinct members of the gamma as
	well as the alhfa and beta subunits, and that preferential associations between members of
	subunit families increase G protein functional diversity.
Gene ID:	2787
Pathways:	Myometrial Relaxation and Contraction, SARS-CoV-2 Protein Interactome
Application Details	
Application Notes:	IF(IHC-P) 1:50-200
	IF(IHC-F) 1:50-200
	IF(ICC) 1:50-200
Restrictions:	For Research Use only
Handling	
Format:	Liquid
Concentration:	1 μg/μL
Buffer:	Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and
	50 % Glycerol.
Preservative:	ProClin
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be
	handled by trained staff only.
Storage:	-20 °C

Handling

Storage Comment:	Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.
Expiry Date:	12 months